arXiv:0902.4732v3 [math.NT] 6 Sep 2009

On the number $\zeta(3)$

L.A.Gutnik

To the thirtieth anniversary
of Apery's discovery.

Table of contents

§0. Foreword.
§1. Short proof of Yu.V. Nesterenko expansion.
§2. Introduction. Begin of the proof of Theorem A.
§3. Transformation the system considered in the Itroduction in the case $\alpha = 1$.
§4. Calculation of the matrix $A^{**}_{1,0}(z, \nu)$.
§5. Further properties of the matrix $A^{*0*}_{1,0}(z, \nu)$.
§6. Further properties of the functions considered in the Introduction.
§7. End of the proof of theorem A.

§0. Foreword .

Let is given a difference equation

$$x_{\nu+1} - b_{\nu+1}x_\nu - a_{\nu+1}x_{\nu-1} = 0, \tag{1}$$

with $\nu \in \mathbb{N}_0$. We denote by

$$\{P_\nu(b_0,\, a_1,\, b_1,\, ...,\, a_\nu,\, b_\nu)\}_{\nu=-1}^{+\infty}$$

and

$$\{Q_\nu(b_0,\, a_1,\, b_1,\, ...,\, a_\nu,\, b_\nu)\}_{\nu=-1}^{+\infty}$$

the solutions of this equation with initial values

$$P_{-1} = 1,\, Q_{-1} = 0,\, P_0(b_0) = b_0,\, Q_0(b_0) = 1. \tag{2}$$

Then

$$\left\{\frac{P_\nu(b_0,\, a_1,\, b_1,\, ...,\, a_\nu,\, b_\nu)}{Q_\nu(b_0,\, a_1,\, b_1,\, ...,\, a_\nu,\, b_\nu)}\right\}_{\nu=0}^{+\infty}$$

is sequence of convergents of continuous fraction

$$b_0+\frac{a_1|}{|b_1}+...+\frac{a_\nu|}{|b_\nu}+... \,.$$

Accoding to the famous result of R. Apéry [1],

$$\zeta(3)=\lim_{\nu\to\infty}\frac{v_\nu}{u_\nu}, \tag{3}$$

where $\{u_\nu\}_{\nu=1}^{+\infty}$ and $\{v_\nu\}_{\nu=1}^{+\infty}$ are solutions of difference equation

$$(\nu+1)^3x_{\nu+1}-(34\nu^3+51\nu^2+27\nu+5)x_\nu+\nu^3x_{\nu-1}=0, \tag{4}$$

with initial values $u_0=1$, $u_1=5$, $v_1=0$, $v_1=6$. The equality (3) is equivalent to the equality

$$\zeta(3)=b^\vee_0+\frac{a^\vee_1|}{|b^\vee_1}+\frac{a^\vee_2|}{|b^\vee_2}+...+\frac{a^\vee_\nu|}{|b^\vee_\nu}+... \tag{5}$$

, with

$$b^\vee_0=0,\ b^\vee_1=5,\ a^\vee_1=6,\ b^\vee_{\nu+1}= \tag{6}$$

$$34\nu^3+51\nu^2+27\nu+5,\ a^\vee_{\nu+1}=-\nu^6,$$

where $\nu\in\mathbb{N}$. Yu.V. Nsterenko in [3] has offered the following expansion the number $2\zeta(3)$ in continuous fraction:

$$2\zeta(3)=2+\frac{1|}{|2}+\frac{2|}{|4}+\frac{1|}{|3}+\frac{4|}{|2}..., \tag{7}$$

with

$$b_0=b_1=a_2=2,\ a_1=1,\ b_2=4, \tag{8}$$

$$b_{4k+1}=2k+2,\ a_{4k+1}=k(k+1),\ b_{4k+2}= \tag{9}$$

$$2k+4,\ a_{4k+2}=(k+1)(k+2)$$

for $k\in\mathbb{N}$,

$$b_{4k+3}=2k+3,\ a_{4k+3}=(k+1)^2,\ b_{4k+4}= \tag{10}$$

$$2k+2,\ a_{4k+4}=(k+2)^2$$

for $k\in\mathbb{N}_0$. The halfs of convergents of continuous fraction (7) compose a sequence containig convergents of continuous fraction (5).

Making use of the method developed in our papers [7] – [17], we have found the followng expansions of the Number $\zeta(3)$ in contiuous fractions :

Teorem A. *The followng equalities hold*

$$(11)\qquad 2\zeta(3)=b_0^{(*1)}+\frac{a_1^{(*1)}|}{|b_1^{(*1)}}+...+\frac{a_\nu^{(*1)}|}{|b_\nu^{(*1)}}+...,$$

$$(12)\qquad 2\zeta(3)=b_0^{(*2)}+\frac{a_1^{(*2)}|}{|b_2^{(*2)}}+...+\frac{a_\nu^{(*2)}|}{|b_\nu^{(*2)}}+...,$$

with

$$b_0^{(*1)}=3,\ a_1^{(*1)}=-81,$$

$$a_\nu^{(*1)}=-(\nu-1)^3\nu^3(4\nu^2-4\nu-3)^3$$

for $\nu\in[2,+\infty)\cap\mathbb{N}$,

$$b_\nu^{(*1)}=4(68\nu^6-45\nu^4+12\nu^2-1)$$

for $\nu\in\mathbb{N}$,

$$b_0^{(*2)}=2,\ a_1^{(*2)}=42,$$

$$a_\nu^{(*2)}=-(\nu-1)^3\nu^3(4\nu^2-4\nu-3)((\nu+1)^3-\nu^3)((\nu-1)^3-(\nu-2)^3)$$

for $\nu\in[2,+\infty)\cap\mathbb{N}$,

$$b_\nu^{(*2)}=2(102\nu^6-68\nu^4+21\nu^2-3),$$

for $\nu\in\mathbb{N}$. I give short proof of Yu.V. Nesterenko expansion in section 1. I prove Theorem A in sections 2 – 7.

§1. Short proof of Yu.V. Nesterenko expansion.

Instead of expansion (7) with (8), it is more convenient for us to prove the equivalent expansion

$$(13)\qquad \zeta(3)=1+\frac{1|}{|4}+\frac{4|}{|4}+\frac{1|}{|3}+\frac{4|}{|2}...,$$

with

$$(14)\qquad b_0=1,\ a_1=1,\ b_1=a_2=b_2=4.$$

Furthermore, to avoid of mishmash in notations, we denote below a_ν, b_ν for the fraction (13) by $a_\nu^\wedge$, $b_\nu^\wedge$. Let $P_{-1}^\vee=1$, $Q_{-1}^\vee=0$,

$$P_\nu^\vee=P_\nu(b_0^\vee,\ a_1^\vee,\ b_1^\vee,\ ...,\ a_\nu^\vee,\ b_\nu^\vee),\ Q_\nu^\vee=Q_\nu(b_0^\vee,\ a_1^\vee,\ b_1^\vee,\ ...,\ a_\nu^\vee,\ b_\nu^\vee),$$

where values $a_\nu^\vee$, $b_\nu^\vee$ are spcified in (6), and $\nu\in\mathbb{N}_0$. Then

$$(15)\qquad Q_0^\vee=1,\ P_0^\vee=b_0^\vee=0,\ Q_1^\vee=b_1^\vee=5,\ P_1^\vee=$$

$$a_1^\vee=6,\ b_2^\vee=117,\ a_2^\vee=-1, P_2^\vee=$$

$$b_2^\vee P_1^\vee + a_2^\vee P_0^\vee = 702, Q_2^\vee =$$
$$b_2^\vee Q_1^\vee + a_2^\vee Q_0^\vee = 584.$$

Let $P_{-1}^\wedge = 1$, $Q_{-1}^\wedge = 0$,

$$P_\nu^\wedge = P_\nu(b_0^\wedge,\, a_1^\wedge,\, b_1^\wedge,\, ...,\, a_\nu^\wedge,\, b_\nu^\wedge),\ Q_\nu^\wedge = Q_\nu(b_0^\wedge,\, a_1^\wedge,\, b_1^\wedge,\, ...,\, a_\nu^\wedge,\, b_\nu^\wedge),$$

where $\nu \in \mathbb{N}_0$, $a_\nu^\wedge := a_\nu$, $b_\nu^\wedge := b_\nu$, and values a_ν, b_ν are spcified in (14), (9), and (10). Then, since $P_{-1}^\wedge = 1$, $Q_{-1}^\wedge = 0$, it follows from (14) that

$$P_0^\wedge = b_0 = 1,\, Q_0^\wedge = 1,\, P_1^\wedge = b_1^\wedge P_0^\wedge + a_1^\wedge P_{-1}^\wedge = 5,\, Q_1^\wedge = b_1^\wedge Q_0^\wedge + a_1^\wedge Q_{-1}^\wedge = 4,$$

$$P_2^\wedge = b_2^\wedge P_1^\wedge + a_2^\wedge P_0^\wedge = 24 = 4P_1^\vee,\, Q_2^\wedge = b_2^\wedge Q_1 + \tag{16}$$
$$a_2^\wedge Q_0 = 20 = 4Q_1^\vee,\, P_3^\wedge = b_3^\wedge P_2^\wedge + a_3^\wedge P_1^\wedge = 77,\, Q_3^\wedge = b_3^\wedge Q_2^\wedge +$$
$$a_3^\wedge Q_1^\wedge = 64,\, P_4^\wedge = b_4^\wedge P_3^\wedge + a_4^\wedge P_2^\wedge = 250,\, Q_4^\wedge = b_4^\wedge Q_3^\wedge + a_4^\wedge Q_2^\wedge = 208,$$
$$P_5^\wedge = b_5^\wedge P_4^\wedge + a_5^\wedge P_3^\wedge = 1154,\, Q_5^\wedge = b_5^\wedge Q_4^\wedge + a_5^\wedge Q_3^\wedge = 960,$$

$$P_6^\wedge = b_6^\wedge P_5^\wedge + a_6^\wedge P_4^\wedge = 12 \times 702 = 12P_2^\vee,\, Q_6^\wedge = b_6^\wedge Q_5^\wedge + \tag{17}$$
$$a_6^\wedge Q_4^\wedge = 12 \times 584 = 12Q_2^\vee.$$

Let

$$U_\nu^\wedge = \begin{pmatrix} P_{\nu-1}^\wedge & Q_{\nu-1}^\wedge \\ P_\nu^\wedge & Q_\nu^\wedge \end{pmatrix}, A_\nu^\wedge = \begin{pmatrix} 0 & 1 \\ a_{1+\nu}^\wedge & b_{1+\nu}^\wedge \end{pmatrix}, \tag{18}$$

$$U_\nu^\vee = \begin{pmatrix} P_{\nu-1}^\vee & Q_{\nu-1}^\vee \\ P_\nu^\vee & Q_\nu^\vee \end{pmatrix}, A_\nu^\vee = \begin{pmatrix} 0 & 1 \\ a_{1+\nu}^\vee & b_{1+\nu}^\vee \end{pmatrix}, \tag{19}$$

where $\nu \in \mathbb{N}_0$. Then

$$U_\nu^\wedge = A_{\nu-1}^\wedge U_{\nu-1}^\wedge,\, U_\nu^\vee = A_{\nu-1}^\vee U_{\nu-1}^\vee \tag{20}$$

for $\nu \in \mathbb{N}$,

$$U_0^\vee = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, U_1^\vee = \begin{pmatrix} 0 & 1 \\ 6 & 5 \end{pmatrix}, U_2^\vee = \begin{pmatrix} 6 & 5 \\ 702 & 584 \end{pmatrix}, \tag{21}$$

$$U_0^\wedge = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, U_1^\wedge = \begin{pmatrix} 1 & 1 \\ 5 & 4 \end{pmatrix}, U_2^\wedge = \begin{pmatrix} 5 & 4 \\ 24 & 20 \end{pmatrix}, U_3^\wedge = \tag{22}$$
$$\begin{pmatrix} 24 & 20 \\ 77 & 64 \end{pmatrix}, U_4^\wedge = \begin{pmatrix} 77 & 64 \\ 250 & 208 \end{pmatrix}, U_5^\wedge =$$
$$\begin{pmatrix} 250 & 208 \\ 1154 & 960 \end{pmatrix}, U_6^\wedge = \begin{pmatrix} 1154 & 960 \\ 8424 & 7008 \end{pmatrix},$$

$$(U_1^{\vee})(U_2^{\wedge})^{-1}=\frac{1}{4}\begin{pmatrix}-24 & 5\\ 0 & 1\end{pmatrix}, \tag{23}$$

$$(U_2^{\vee})(U_6^{\wedge})^{-1}=\frac{1}{96}\begin{pmatrix}-36 & 5\\ 0 & 8\end{pmatrix}. \tag{24}$$

Let

$$H_1=\frac{1}{4}\begin{pmatrix}-24 & 5\\ 0 & 1\end{pmatrix}, \tag{25}$$

$$H_k=\begin{pmatrix}12(k+2)(k+1)c(k+1) & -5(k+2)c(k+1)\\ 0 & -(k-1)^3c(k)\end{pmatrix}, \tag{26}$$

where $k\in[2,+\infty)\cap\mathbb{Z}$ and $c(k)=(-2(k-1)^3(k+1)!)^{-1}$. Let $k\in\mathbb{N}$, $k\geq 2$. Then, in view of (18),

$$A_{4k-6}^{\wedge}=\begin{pmatrix}0 & 1\\ a_{4(k-2)+3}^{\wedge} & b_{4(k-2)+3}^{\wedge}\end{pmatrix}=\begin{pmatrix}0 & 1\\ (k-1)^2 & 2k-1\end{pmatrix},$$

$$A_{4k-5}^{\wedge}=\begin{pmatrix}0 & 1\\ a_{4(k-2)+4}^{\wedge} & b_{4(k-2)+4}^{\wedge}\end{pmatrix}=\begin{pmatrix}0 & 1\\ k^2 & 2k-2\end{pmatrix},$$

$$A_{4k-4}^{\wedge}=\begin{pmatrix}0 & 1\\ a_{4(k-1)+1}^{\wedge} & b_{4(k-1)+1}^{\wedge}\end{pmatrix}=\begin{pmatrix}0 & 1\\ k^2-k & 2k\end{pmatrix},$$

$$A_{4k-3}^{\wedge}=\begin{pmatrix}0 & 1\\ a_{4(k-1)+2}^{\wedge} & b_{4(k-1)+2}^{\wedge}\end{pmatrix}=\begin{pmatrix}0 & 1\\ k^2+k & 2k+2\end{pmatrix},$$

$$A_{4k-5}^{\wedge}A_{4k-6}^{\wedge}=\begin{pmatrix}(k-1)^2 & 2k-1\\ 2(k-1)^3 & 5k^2-6k+2\end{pmatrix},$$

$$A_{4k-4}^{\wedge}A_{4k-5}^{\wedge}A_{4k-6}^{\wedge}=\begin{pmatrix}2(k-1)^3 & 5k^2-6k+2\\ 5k(k-1)^3 & k(12k^2-15k+5)\end{pmatrix}.$$

Let

$$B_k^{\wedge}=A_{4k-3}^{\wedge}A_{4k-4}^{\wedge}A_{4k-5}^{\wedge}A_{4k-6}= \tag{27}$$

$$\begin{pmatrix}5k(k-1)^3 & k(12k^2-15k+5)\\ 12k(k+1)(k-1)^3 & k(k+1)(29k^2-36k+12)\end{pmatrix},$$

Then, in view of (20),

$$U_{4k-2}^{\wedge}=B_k^{\wedge}U_{4k-6}^{\wedge}, \tag{28}$$

where, as before, $k\in[2,+\infty)\cap\mathbb{Z}$. Let now $k\in[3,+\infty)\cap\mathbb{Z}$. Then, in view of (26),

$$H_{k-1}=\begin{pmatrix}12(k+1)kc(k) & -5(k+1)c(k)\\ 0 & -(k-2)^3c(k-1)\end{pmatrix}. \tag{29}$$

In view of (6)

$$b_k^{\vee} = 34(k-1)^3 + 51(k-1)^2 + 27(k-1) + 5 = \tag{30}$$

$$34k^3 - 51k^2 + 27k - 5,\ a_k^{\vee} = -(k-1)^6,$$

where $k \in [3, +\infty) \cap \mathbb{Z}$. Hence, in view of (19),

$$A_{k-1}^{\vee} = \begin{pmatrix} 0 & 1 \\ -(k-1)^6 & 34k^3 - 51k^2 + 27k - 5 \end{pmatrix}, \tag{31}$$

In view (29) – (31),

$$A_{k-1}^{\vee} H_{k-1} = \begin{pmatrix} 0 & 1 \\ -(k-1)^6 & 34k^3 - 51k^2 + 27k - 5 \end{pmatrix} \times \tag{32}$$

$$\begin{pmatrix} 12(k+1)kc(k) & -5(k+1)c(k) \\ 0 & -(k-2)^3 c(k-1) \end{pmatrix} =$$

$$\begin{pmatrix} 0 & -(k-2)^3 c(k-1) \\ -(k-1)^6 12(k+1)kc(k) & (k-1)^6 5(k+1)c(k) - b_k^{\vee}(k-2)^6 c(k-1). \end{pmatrix}.$$

In view (27) and (26),

$$H_k B_k^{\wedge} = \begin{pmatrix} 12(k+2)(k+1)c(k+1) & -5(k+2)c(k+1) \\ 0 & -(k-1)^3 c(k) \end{pmatrix} \times \tag{33}$$

$$\begin{pmatrix} 5k(k-1)^3 & k(12k^2 - 15k + 5) \\ 12k(k+1)(k-1)^3 & k(k+1)(29k^2 - 36k + 12) \end{pmatrix} =$$

$$\begin{pmatrix} 0 & (k+2)c(k+1)k(k+1)(-k^2) \\ -c(k)12k(k+1)(k-1)^6 & -(k-1)^3 c(k)k(k+1)(29k^2 - 36k + 12) \end{pmatrix}.$$

Since

$$-(k+2)(k+1)c(k+1)k^3 = -c(k-1)(k-2)^3,$$

$$-(k-1)^3 c(k)k(k+1)(29k^2 - 36k + 12) - ((k-1)^6 5(k+1)c(k) =$$

$$-(34k^3 - 51k^2 + 27k - 5)(k-1)^3(k+1)c(k)) =$$

$$-(34k^3 - 51k^2 + 27k - 5)(k-2)^3)c(k-1)),$$

it follows from (30), (32) and (33) that

$$A_{k-1}^{\vee} H_{k-1} = H_k B_k^{\wedge} \tag{34}$$

for $k \in [3, +\infty) \cap \mathbb{Z}$. We prove by induction now the followiong equality

$$U_k^{\vee} = H_k U_{4k-2}^{\wedge}. \tag{35}$$

for any $k \in \mathbb{N}$. In view of (23) and (25), the equality (35) holds for $k = 1$. In view of (24) and (26), the equality (35) holds for $k = 2$. Let $k \in [3, +\infty) \cap \mathbb{Z}$ and (35) holds for $k - 1$. Then, in view of(28), (34) and (20),

$$H_k U_{4k-2}^{\wedge} = H_k B_k U_{4k-6}^{\wedge} = A_{k-1}^{\vee} H_{k-1} U_{4k-6}^{\wedge} = A_{k-1}^{\vee} U_{k-1}^{\vee} = U_k^{\vee}.$$

So, the equality (35) holds for any $k \in \mathbb{N}$. In view of (35),

$$P^{\vee}_{k} = (2(k+1)!)^{-1}P^{\wedge}_{4k-2},\ Q^{\vee}_{k} = (2(k+1)!)^{-1}Q^{\wedge}_{4k-2} \tag{36}$$

for $k \in [2, +\infty) \cap \mathbb{Z}$. Since

$$P^{\vee}_{\nu} = (\nu!)^3 v_\nu,\ Q^{\vee}_{\nu} = (\nu!)^3 u_\nu \tag{37}$$

for v_ν and u_ν in (3) and $\nu \in \mathbb{N}_0$, it follows from (36) and (37), that

$$P^{\wedge}_{4k-2} = 2(k+1)(k!)^4 v_k,\ Q^{\wedge}_{4k-2} = 2(k+1)(k!)^4 u_k \tag{38}$$

Let $k \in [2, +\infty) \cap \mathbb{Z}$. In view of (14) and (9) - (10),

$$Q^{\wedge}_{4k-2} < Q^{\wedge}_{4k-1} < Q^{\wedge}_{4k-2}(k^2+2k+1), \tag{39}$$

$$\begin{gathered} Q^{\wedge}_{4k-2} < Q^{\wedge}_{4k-1} < Q^{\wedge}_{4k} < \\ Q^{\wedge}_{4k-1}(k^2+4k+1) < Q^{\wedge}_{4k-2}(k^2+2k+1)(k^2+4k+1), \end{gathered} \tag{40}$$

$$\begin{gathered} Q^{\wedge}_{4k-2} < Q^{\wedge}_{4k} < Q^{\wedge}_{4k+1} < Q^{\wedge}_{4k}(k^2+3k+2) < \\ Q^{\wedge}_{4k-2}(k^2+2k+1)(k^2+4k+1)(k^2+3k+2), \end{gathered} \tag{41}$$

$$\begin{gathered} \prod_{\kappa=1}^{4k-2} a_\kappa = 4\prod_{\kappa=2}^{k} a_{4\kappa-5}a_{4\kappa-4}a_{4\kappa-3}a_{4\kappa-2} = \\ 4\prod_{\kappa=2}^{k}(\kappa-1)^2\kappa^2(\kappa-1)\kappa\kappa(\kappa+1) = 2(k!)^8(k+1)/k^3, \end{gathered} \tag{42}$$

$$\prod_{\kappa=1}^{4k-1} a_\kappa = k^2\prod_{\kappa=1}^{4k-2} a_\kappa = 2(k!)^8(k+1)/k, \tag{43}$$

$$\prod_{\kappa=1}^{4k} a_\kappa = (k+1)^2\prod_{\kappa=1}^{4k-1} a_\kappa = 2(k!)^8(k+1)^3/k, \tag{44}$$

$$\prod_{\kappa=1}^{4k+1} a_\kappa = k(k+1)\prod_{\kappa=1}^{4k} a_\kappa = 2(k!)^8(k+1)^4. \tag{45}$$

As it is well known, for any $\varepsilon > 0$ there exist $C_1(\varepsilon) > 0$ and $C_2(\varepsilon) > 0$ such that

$$C_1(\varepsilon)(1+\sqrt{2})^{4k(1-\varepsilon)} < |u_k| < C_2(\varepsilon)(1+\sqrt{2})^{4k(1+\varepsilon)}, \tag{46}$$

(47) $$C_1(\varepsilon)(1+\sqrt{2})^{4k(1-\varepsilon)} < |v_k| < C_2(\varepsilon)(1+\sqrt{2})^{4k(1+\varepsilon)},$$

(48) $$\frac{C_1(\varepsilon)}{(1+\sqrt{2})^{8k(1+\varepsilon)}} < \left|\zeta(3)-\frac{v_k}{u_k}\right| < \frac{C_2(\varepsilon)}{(1+\sqrt{2})^{8k(1-\varepsilon)}}.$$

Terefore, according to (38) – (48), and well known expression for the difference of two neighboring convergents of continuous fraction, for any $\varepsilon > 0$ there exist $C_3(\varepsilon) > 0$ and $C_4(\varepsilon) > 0$ such that

(49) $$\frac{C_3(\varepsilon)}{(1+\sqrt{2})^{8k(1+\varepsilon)}} < \left|\zeta(3)-\frac{P^{\wedge}_{4k-2}}{Q^{\wedge}_{4k-2}}\right| < \frac{C_4(\varepsilon)}{(1+\sqrt{2})^{8k(1-\varepsilon)}},$$

(50) $$\frac{C_3(\varepsilon)}{(1+\sqrt{2})^{8k(1+\varepsilon)}} < \left|\frac{P^{\wedge}_{4k-1+i}}{Q^{\wedge}_{4k-1+i}}-\frac{P^{\wedge}_{4k-2+i}}{Q^{\wedge}_{4k-2+i}}\right| < \frac{C_4(\varepsilon))}{(1+\sqrt{2})^{8k(1-\varepsilon)}}$$

for $i = 0,\ 1,\ 2$, and

(51) $$\frac{C_3(\varepsilon)}{(1+\sqrt{2})^{8k(1-\varepsilon)}} < \left|\zeta(3)-\frac{P^{\wedge}_{4k-2+i}}{Q^{\wedge}_{4k-2+i}}\right| < \frac{C_4(\varepsilon)}{(1+\sqrt{2})^{8k(1+\varepsilon)}},$$

for $i = 0,\ 1,\ 2,\ 3$. ■.

§2. Introduction. Begin of the proof of Theorem A.

Let

(52) $$|z| > 1, -3\pi/2 < \arg(z) \le \pi/2, \log(z) = \ln(|z|) + i\arg(z).$$

Then $\log(-z) = \log(z) - i\pi$, if $\Re(z) > 0$ and $\log(z) = \log(-z) - i\pi$, if $\Re(z) < 0$. Let $\alpha \in \mathbb{N}_0$,

(53) $$f^{*\vee}_{\alpha,0,1}(z,\nu) := f^{*}_{\alpha,0,1}(z,\nu) := \sum_{k=0}^{\nu+\alpha}(z)^k\binom{\nu+\alpha}{k}^2\binom{\nu+k}{\nu}^2,$$

(54) $$R(\alpha,t,\nu) = \frac{\prod_{j=1}^{\nu}(t-j)}{\prod_{j=0}^{\nu+\alpha}(t+j)},$$

(55) $$f^{*}_{\alpha,0,2}(z,\nu) = \sum_{t=1}^{+\infty} z^{-t}\frac{(\nu+\alpha)!^2}{\nu!^2}(R(\alpha,t,\nu))^2,$$

(56) $$f^{*}_{\alpha,0,4}(z,\nu) = -\sum_{t=1}^{+\infty} z^{-t}\frac{(\nu+\alpha)!^2}{\nu!^2}\left(\frac{\partial}{\partial t}(R^2)\right)(\alpha,t,\nu),$$

$$f^*_{\alpha,0,3}(z,\nu)=(\log(z))f^*_{\alpha,0,2}(z,\nu)+f^*_{\alpha,0,4}(z,\nu), \tag{57}$$

$$f_{\alpha,0,k}=\frac{\nu!^2}{(\nu+\alpha)!^2}(z,\nu)f^*_{\alpha,0,k}(z,\nu) \tag{58}$$

where $k=1,\ 2,\ 3,\ 4,\ \nu\in\mathbb{N}_0$. Let

$$\mu=\mu_\alpha(\nu)=(\nu+\alpha)(\nu+1),\ \tau=\tau_\alpha(\nu)=\nu+\frac{1+\alpha}{2}, \tag{59}$$

$$\begin{gathered}a^{\vee}_{\alpha,0,1,1}(z,\nu)=\frac{1}{2}(-1+2\alpha-\alpha^2-5\mu+3\alpha\mu-5\mu^2-\alpha\mu^2)+\\ \frac{z}{2}(-4+12\alpha-13\alpha^2+6\alpha^3-\alpha^4)+\\ \frac{z}{2}\mu(-32+54\alpha-29\alpha^2+5\alpha^3-56\mu+20\alpha\mu),\end{gathered} \tag{60}$$

$$\begin{gathered}a^{\vee}_{\alpha,0,1,2}(z;\nu)=-2+3\alpha-\alpha^2-8\mu+\alpha\mu-\alpha^2\mu-4\mu^2+\\ z(2-11\alpha+17\alpha^2-10\alpha^3+2\alpha^4-4\mu-11\alpha\mu+3\alpha^2\mu-20\mu^2),\end{gathered} \tag{61}$$

$$\begin{gathered}a^{\vee}_{\alpha,0,1,3}(z;\nu)=-4+5\alpha-\frac{3}{2}\alpha^2-\frac{1}{2}\alpha^3-12\mu-2\alpha\mu+\\ z\left(10-24\alpha+\frac{37}{2}\alpha^2-\frac{11}{2}\alpha^3+24\mu-22\alpha\mu\right),\end{gathered} \tag{62}$$

$$a^{\vee}_{\alpha,0,1,4}(z;\nu)=(z-1)(6-7\alpha+3\alpha^2+12\mu), \tag{63}$$

$$\begin{gathered}a^{\wedge}_{\alpha,0,1,1}(z;\nu)=1-\alpha+3\mu+\mu^2+\\ z(4-8\alpha+5\alpha^2-\alpha^3+24\mu-22\alpha\mu+5\alpha^2\mu+16\mu^2),\end{gathered} \tag{64}$$

$$\begin{gathered}a^{\wedge}_{\alpha,0,1,2}(z;\nu)=4-2\alpha+8\mu+2\alpha\mu+\\ z(-4+18\alpha-16\alpha^2+4\alpha^3+16\mu+10\alpha\mu),\end{gathered} \tag{65}$$

$$\begin{gathered}a^{\wedge}_{\alpha,0,1,3}(z;\nu)=\\ 8-2\alpha+\alpha^2+8\mu+z(-20+28\alpha-5\alpha^2-8\mu),\end{gathered} \tag{66}$$

$$a^{\wedge}_{\alpha,0,1,4}(z;\nu)=-(z-1)(12-2\alpha), \tag{67}$$

$$a^{\vee}_{\alpha,0,2,1}(z;\nu)=\frac{z}{2}(-4+12\alpha-13\alpha^2+6\alpha^3-\alpha^4)+ \tag{68}$$

$$\frac{z}{2}\mu(-32+54\alpha-29\alpha^2+5\alpha^3)+\frac{z}{2}\mu^2(-68+34\alpha-6\alpha^2-24\mu),$$

$$a^{\vee}_{\alpha,0,2,2}(z;\nu)=\frac{1}{2}(-1+2\alpha-\alpha^2-5\mu+3\alpha\mu-5\mu^2-\alpha\mu^2)+ \tag{69}$$
$$\frac{z}{2}(-10\alpha+21\alpha^2-14\alpha^3+3\alpha^4)+$$
$$\frac{z}{2}\mu(-16-20\alpha+17\alpha^2-7\alpha^3-48\mu-28\alpha\mu),$$

$$a^{\vee}_{\alpha,0,2,3}(z;\nu)=-2+3\alpha-\alpha^2-8\mu+\alpha\mu-\alpha^2\mu-4\mu^2+ \tag{70}$$
$$\frac{z}{2}(12-32\alpha+25\alpha^2-5\alpha^3-2\alpha^4)+z\mu(20-23\alpha-3\alpha^2+4\mu),$$

$$a^{\vee}_{\alpha,0,2,4}(z;\nu)=\frac{1}{2}(-8+10\alpha-3\alpha^2-\alpha^3-24\mu-4\alpha\mu)+ \tag{71}$$
$$\frac{z}{2}(8-10\alpha+3\alpha^2+\alpha^3+24\mu+4\alpha\mu),$$

$$a^{\vee}_{\alpha,0,3,1}(z;\nu)=\frac{z}{2}(-4+12\alpha-13\alpha^2+6\alpha^3-\alpha^4)+ \tag{72}$$
$$\frac{z}{2}\mu(-32+54\alpha-29\alpha^2+5\alpha^3)+$$
$$\frac{z}{2}\mu^2(-76+44\alpha-9\alpha^2-\alpha^3-48\mu-4\alpha\mu),$$

$$a^{\vee}_{\alpha,0,3,2}(z;\nu)=z(-2+\alpha+4\alpha^2-4\alpha^3+\alpha^4)+ \tag{73}$$
$$z\mu(-16-\alpha+7\alpha^2-3\alpha^3-\alpha^4-34\mu-17\alpha\mu-7\alpha^2\mu-12\mu^2),$$

$$a^{\vee}_{\alpha,0,3,3}(z;\nu)=\frac{1}{2}(-1+2\alpha-\alpha^2-5\mu+3\alpha\mu-5\mu^2-\alpha\mu^2)+ \tag{74}$$
$$\frac{z}{2}(4-16\alpha+15\alpha^2-4\alpha^3-\alpha^5)+$$
$$\frac{z}{2}\mu(16-42\alpha+\alpha^2-9\alpha^3+8\mu-20\alpha\mu),$$

$$a^{\vee}_{\alpha,0,3,4}(z;\nu)=-2+3\alpha-\alpha^2-8\mu+\alpha\mu-\alpha^2\mu-4\mu^2+ \tag{75}$$
$$z(2-3\alpha+\alpha^2+8\mu-\alpha\mu+\alpha^2\mu+4\mu^2),$$

$$a^{\vee}_{\alpha,0,4,1}(z;\nu)=\frac{z}{2}(-4+12\alpha-13\alpha^2+6\alpha^3-\alpha^4)+ \tag{76}$$
$$\frac{z}{2}\mu(-32+54\alpha-29\alpha^2+5\alpha^3-80\mu+50\alpha\mu-11\alpha^2\mu-\alpha^3\mu)+$$

$$z\mu^3(-32-\alpha-\alpha^2-4\mu),$$

$$a^{\vee}_{\alpha,0,4,2}(z;\nu)=\frac{z}{2}(-8+14\alpha-5\alpha^2-2\alpha^3+\alpha^4)+ \tag{77}$$

$$\frac{z}{2}\mu(-56+32\alpha+\alpha^2-5\alpha^3-2\alpha^4)+$$

$$\frac{z}{2}\mu^2(-112-26\alpha-15\alpha^2-5\alpha^3)+z\mu^3(-28-10\alpha),$$

$$a^{\vee}_{\alpha,0,4,3}(z;\nu)=\frac{z}{2}(-4+5\alpha^2-2\alpha^3-\alpha^5)+ \tag{78}$$

$$\frac{z}{2}\mu(-24-22\alpha-3\alpha^2-9\alpha^3-4\alpha^4-36\mu-42\alpha\mu-18\alpha^2\mu-8\mu^2),$$

$$a^{\vee}_{\alpha,0,4,4}(z;\nu)=\frac{1}{2}(-1+2\alpha-\alpha^2-5\mu+3\alpha\mu-5\mu^2-\alpha\mu^2)+ \tag{79}$$

$$\frac{z}{2}(-2\alpha+\alpha^2-\alpha^5)+\frac{z}{2}\mu(-8\alpha-5\alpha^2-5\alpha^3-4\alpha\mu),$$

$$a^{\wedge}_{\alpha,0,2,1}(z;\nu)=z(4-8\alpha+5\alpha^2-\alpha^3)+ \tag{80}$$

$$z\mu(24-22\alpha+5\alpha^2+28\mu-2\alpha\mu),$$

$$a^{\wedge}_{\alpha,0,2,2}(z;\nu)=1-\alpha+3\mu+\mu^2+ \tag{81}$$

$$z(10\alpha-11\alpha^2+3\alpha^3)+z\mu(16+16\alpha+\alpha^2)+16z\mu^2,$$

$$a^{\wedge}_{\alpha,0,2,3}(z;\nu)=4-2\alpha+8\mu+2\alpha\mu+ \tag{82}$$

$$z(-12+20\alpha-5\alpha^2+2\alpha^3)+z\mu(-16+14\alpha),$$

$$a^{\wedge}_{\alpha,0,2,4}(z;\nu)=(8-2\alpha+\alpha^2+8\mu)(1-z), \tag{83}$$

$$a^{\wedge}_{\alpha,0,3,1}(z;\nu)=z(4-8\alpha+5\alpha^2-\alpha^3)+ \tag{84}$$

$$z\mu(24-22\alpha+5\alpha^2+36\mu-4\alpha\mu+\alpha^2\mu+8\mu^2),$$

$$a^{\wedge}_{\alpha,0,3,2}(z;\nu)=z(4+2\alpha-6\alpha^2+2\alpha^3)+ \tag{85}$$

$$z\mu(24+14\alpha+2\alpha^3)+z\mu^2(28+14\alpha),$$

$$a^{\wedge}_{\alpha,0,3,3}(z;\nu)=1-\alpha+3\mu+\mu^2+ \tag{86}$$

$$z(-4+12\alpha-3\alpha^2+\alpha^3+\alpha^4-8\mu+18\mu\alpha+7\mu\alpha^2),$$

(87) $$a^{\wedge}_{\alpha,0,3,4}(z;\nu)=(4-2\alpha+8\mu+2\alpha\mu)(1-z),$$

(88) $$a^{\wedge}_{\alpha,0,4,1}(z;\nu)=z(4-8\alpha+5\alpha^2-\alpha^3)+$$
$$z\mu(24-22\alpha+5\alpha^2+40\mu-6\alpha\mu+\alpha^2\mu)+z\mu^3(16+2\alpha),$$

(89) $$a^{\wedge}_{\alpha,0,4,2}(z;\nu)=z(8-6\alpha-\alpha^2+\alpha^3)+$$
$$z\mu(40+4\alpha+\alpha^2+2\alpha^3)+z\mu^2(48+22\alpha+5\alpha^2)+8z\mu^3,$$

(90) $$a^{\wedge}_{\alpha,0,4,3}(z;\nu)=z(4+4\alpha-\alpha^2+\alpha^3+\alpha^4)+$$
$$z\mu(16+22\alpha+11\alpha^2+4\alpha^3)+z\mu^2(12+10\alpha)$$

(91) $$a^{\wedge}_{\alpha,0,4,4}(z;\nu)=1-\alpha+3\mu+\mu^2+$$
$$2z\alpha+z\alpha^2+z\alpha^3+z\alpha^4+4z\mu\alpha+3z\mu\alpha^2,$$

(92) $$a^{*}_{\alpha,0,i,k}(z;\nu)=a^{\vee}_{\alpha,0,i,k}(z;\nu)+\tau a^{\wedge}_{\alpha,0,i,k}(z;\nu),$$

where $i=1,\ ...,\ 4,\ k=1,\ ...,\ 4$. We denote by

$$A^{*}_{\alpha,0}(z;\nu),\ A^{\vee}_{\alpha,0}(z;\nu)(z;\nu),\ A^{\wedge}_{\alpha,0}(z;\nu)(z;\nu)$$

the 4×4-matrix, such that its element in i-th row and k-th column is equal respectively to

$$a^{*}_{\alpha,0,i,k}(z;\nu),\ a^{\vee}_{\alpha,0,i,k}(z;\nu),\ a^{\wedge}_{\alpha,0,i,k}(z;\nu)$$

where $i=1,\ ...,\ 4,\ k=1,\ ...,\ 4$. Then

(93) $$A^{*}_{\alpha,0}(z;\nu)=A^{\vee}_{\alpha,0}(z;\nu)(z;\nu)+\tau A^{\wedge}_{\alpha,0}(z;\nu)(z;\nu).$$

Clearly,

(94) $$A^{*}_{\alpha,0}(z;\nu)=A^{*}_{\alpha,0}(z;\nu)(1;\nu)+(z-1)V^{*}_{\alpha,0}(\nu),$$

where the matrix $V^{*}_{\alpha,0}(\nu)$ does not depend from z. Let

(95)
$$X_{\alpha,0,k}(z;\nu)=\begin{pmatrix}f_{\alpha,0,k}(z,\nu)\\ \delta f_{\alpha,0,k}(z,\nu)\\ \delta^2 f_{\alpha,0,k}(z,\nu)\\ \delta^3 f_{\alpha,0,k}(z,\nu)\end{pmatrix},\ X^{*}_{\alpha,0,k}(z;\nu)=\frac{((\nu+\alpha)!)^2}{(\nu!)^2}X_{\alpha,0,k}(z;\nu)$$

for $k=1,\ 2,\ 3,\ |z|>1,\ \nu\in\mathbb{N}_0$. Let further

(96) $$X_{\alpha,0,k}(z;-\nu-1-\alpha)=X_{\alpha,0,k}(z;\nu),$$

where $\nu \in \mathbb{N}_0$. The following results are obtained in [15] – [17].

Theorem 1. *The column* $X_{\alpha,0,k}(z;\nu)$ *satisfies to the equation*

$$\nu^5 X_{\alpha,0,k}(z;\nu-1) = A^*_{\alpha,0}(z;\nu)X_{\alpha,0,k}(z;\nu), \tag{97}$$

for $\nu \in M^*_\alpha = (-\infty, -1-\alpha] \cup [1,+\infty)) \cap \mathbb{Z}$, $k = 1,\ 2,\ 3$, $|z| > 1$; *moreover, the matrix* $A^*_{\alpha,0}(z;\nu)$ *has the following property:*

$$-\nu^5(\nu+\alpha)^5 E_4 = A^*_{\alpha,0}(z;-\nu-\alpha)A^*_{\alpha,0}(z;\nu), \tag{98}$$

where E_4 *is the* 4×4 *unit matrix,* $z \in \mathbb{C}$, $\nu \in \mathbb{C}$. Let us consider the row

$$R_{\alpha,0}(\nu) = (r_{\alpha,0,1}(\nu),\ r_{\alpha,0,2}(\nu),\ r_{\alpha,0,3}(\nu),\ r_{\alpha,0,4}(\nu)), \tag{99}$$

where

$$r_{\alpha,0,1}(\nu) = \mu_\alpha(\nu)^2,\ r_{\alpha,0,2}(\nu) = -2(1-\alpha) \times \tag{100}$$

$$\mu_\alpha(\nu),\ r_{\alpha,0,3}(\nu) = (1-\alpha)^2 - 2\mu_\alpha(\nu),\ r_{\alpha,0,4}(0,\nu) = 2(1-\alpha).$$

The following Theorem is proved in [17] (Lemma 11.3.1).

Theorem 2. *The row* $R_{\alpha,0}(\nu)$ *has the following property:*

$$R_{\alpha,0}(\nu-1)A^*_{\alpha,0}(1;\nu) = \nu^5 R_{\alpha,0}(\nu), \tag{101}$$

where $\nu \in \mathbb{C}$.

§3. Transformations of the system considered in the Introduction in the case $\alpha = 1$.

In view of (59), (100)

$$\tau = \tau_1(\nu) = \nu+1,\ \mu = \mu_1(\nu) = (\nu+1)^2,\ r_{1,0,1}(\nu) = \tag{102}$$

$$\mu_1(\nu)^2 = (\nu+1)^4 = \tau^4,\ r_{1,0,2}(\nu) = 0,\ r_{1,0,3}(\nu) =$$

$$-2\mu_1(\nu) = -2(\nu+1)^2,\ r_{\alpha,0,4}(0,\nu) = 0.$$

Let E_4 denotes 4×4-unit matrix, and let $C^{(\kappa)}(\nu)$ with $\kappa = 0,\ 1$ is result of replacement of $2\kappa+1$-th row of the matrix E_4 by the row in (99) with $\alpha = 1$. Let further $D^{(\kappa)}(\nu)$ with $\kappa = 0,\ 1$ denotes the adjoint matrix to the matrix $C^{(\kappa)}(\nu)$. Then

$$C^{(0)}(\nu) = \tag{103}$$

$$\begin{pmatrix} r_{1,0,1}(\nu) & r_{1,0,2}(\nu) & r_{1,0,3}(\nu) & r_{1,0,4}(\nu) \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},\ C^{(1)}(\nu) =$$

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ r_{1,0,1}(\nu) & r_{1,0,2}(\nu) & r_{1,0,3}(\nu) & r_{1,0,4}(\nu) \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

$$D^{(0)}(\nu) = \tag{104}$$

$$\begin{pmatrix} 1 & -r_{1,0,2}(\nu) & -r_{1,0,3}(\nu) & -r_{1,0,4}(\nu) \\ 0 & r_{1,0,1}(\nu) & 0 & 0 \\ 0 & 0 & r_{1,0,1}(\nu) & 0 \\ 0 & 0 & 0 & r_{1,0,1}(\nu) \end{pmatrix},\ D^{(1)}(\nu) =$$

$$\begin{pmatrix} r_{1,0,3}(\nu) & 0 & 0 & 0 \\ 0 & r_{1,0,3}(\nu) & 0 & 0 \\ -r_{1,0,1}(\nu) & -r_{1,0,2}(\nu) & 1 & -r_{1,0,4}(\nu) \\ 0 & 0 & 0 & r_{1,0,3}(\nu) \end{pmatrix}.$$

Clearly,

$$C^{(\kappa)}(\nu)D^{(\kappa)}(\nu) = (-2)^{\kappa}(\mu_1(\nu))^{2-\kappa}E_4,\ C^{(\kappa)}(-\nu-2) = \tag{105}$$

$$C^{(\kappa)}(\nu),\ D^{(\kappa)}(-\nu-2) = D^{(\kappa)}(\nu),$$

where $\kappa = 0,\ 1$. Let

$$A^{*\kappa*}_{1,0}(z,\nu) = C^{(\kappa)}(\nu-1)A^{*}_{1,0}(z,\nu)D^{(\kappa)}(\nu). \tag{106}$$

Then

$$A^{*\kappa*}_{1,0}(z,-\nu-1) = \tag{107}$$

$$C^{(\kappa)}(-\nu-2)A^{*}_{1,0}(z,-\nu-1)D^{(\kappa)}(-\nu-1) =$$

$$C^{(\kappa)}(\nu)A^{*}_{1,0}(z,-\nu-1)D^{(\kappa)}(\nu-1),$$

and, in view of (105), (98), (106),

$$A^{*\kappa*}_{1,0}(z,-\nu-1)A^{*\kappa*}_{1,0}(z,\nu) = \tag{108}$$

$$C^{(\kappa)}(\nu)A^{*}_{1,0}(z,-\nu-1)D^{(\kappa)}(\nu-1)C^{(k)}(\nu-1)A^{*}_{1,0}(z,\nu)D(\nu) =$$

$$-4^{\kappa}(\mu_1(\nu)\mu_1(\nu-1))^{2-\kappa}(\nu(\nu+1))^5E_4,$$

where $\kappa = 0,\ 1$. Let

$$Y^{(\kappa)}_{1,0,k}(z;\nu) = C^{(\kappa)}(\nu)X_{1,0,k}(z;\nu), \tag{109}$$

where $\kappa = 0,\ 1,\ k = 1,\ 2,\ 3,\ |z| > 1,\ \nu \in M^{***}_1 = ((-\infty,-2] \cup [0,+\infty)) \cap \mathbb{Z}$. Then, in view of (96), (105), (97),

$$Y^{(\kappa)}_{1,0,k}(z;-\nu-2) = Y^{(\kappa)}_{1,0,k}(z;\nu), \tag{110}$$

$$A^{*\kappa*}_{1,0}(z,\nu)Y^{(\kappa)}_{1,0,k}(z;\nu) = \tag{111}$$

$$C^{(\kappa)}(\nu-1)A^{*}_{1,0}(z,\nu)D^{(\kappa)}(\nu)C^{(\kappa)}(\nu)X_{1,0,k}(z;\nu) =$$

$$(-1)^{\kappa}(\kappa+1)\mu_1(\nu)^{2-\kappa}C^{(\kappa)}(\nu-1)A^{*}_{1,0}(z,\nu)X_{1,0,k}(z;\nu) =$$

$$(-1)^{\kappa}(\kappa+1)\mu_1(\nu)^{2-\kappa}\nu^5C^{(\kappa)}(\nu-1)X_{1,0,k}(z;\nu-1) =$$

$$(-1)^{\kappa}(\kappa+1)\mu_1(\nu)^{2-\kappa}\nu^5\nu^5Y^{(\kappa)}_{1,0,k}(z;\nu-1),$$

where $\kappa = 0,\ 1$, $k = 1,\ 2,\ 3$, $|z| > 1$, $\nu \in M^{*}_1 = ((-\infty,-2]\cup[1,+\infty))\cap\mathbb{Z}$. Replacing in the equality (111)

$$\nu \in M^{*}_1 = ((-\infty,-2]\cup[1,+\infty))\cap\mathbb{Z}$$

by

$$\nu := -\nu-2 \in M^{**}_1 = ((-\infty,-3]\cup[0,+\infty))\cap\mathbb{Z},$$

and taking in account (110) we obtain the equality

(112)

$$-A^{*\kappa*}_{1,0}(z,-\nu-2)Y^{(\kappa)}_{1,0,k}(z;\nu) = (-1)^{\kappa}(\kappa+1)\mu_1(\nu)^{2-\kappa}(\nu+2)^5Y^{(\kappa)}_{1,0,k}(z;\nu+1),$$

where $\kappa = 0,\ 1$, $k = 1,\ 2,\ 3$, $|z| > 1$, $\nu \in M^{**}_1 = ((-\infty,-3]\cup[0,+\infty))\cap\mathbb{Z}$.

§4. Calculation of the matrix $A^{*\kappa*}_{1,0}(z,\nu)$ for $\kappa = 0,\ 1$.

Let

$$V^{*\kappa*}_{1,0}(\nu) = C^{(\kappa)}(\nu-1)V^{*}_{1,0}(\nu)D^{(\kappa)}(\nu). \tag{113}$$

Then, in view of (94),

$$A^{*\kappa*}_{\alpha,0}(z;\nu) = A^{*\kappa*}_{\alpha,0}(1;\nu)(1;\nu) + (z-1)V^{*\kappa*}_{\alpha,0}(\nu), \tag{114}$$

where the matrix $V^{**}_{\alpha,0}(\nu)$ does not depend from z. We note that the $2\kappa+1$-th row of the matrix

$$C^{(\kappa)}(\nu-1)A^{*}_{1,0}(1,\nu)$$

coincides with the row

$$R^{*}_{1,0}(\nu-1)A^{**}_{1,0}(z,\nu)$$

and, according to the Theorem 2 coincides with the row $\nu^5R^{*}_{1,0}(\nu)$, i.e. with the $2\kappa+1$-th row of the matrix $\nu^5C^{(\kappa)}(\nu)$. Therefore, in view of (105), the $(2\kappa+1)$-th row of the matrix $A^{*\kappa*}_{1,0}(1,\nu)$ is equal to

$$(-2)^{\kappa}(\mu_1(\nu)^{2-\kappa})\nu^5\bar{e}_{4,2\kappa+1}, \tag{115}$$

where $\bar{e}_{4,l}$ denotes the l-th row of the matrix E_4 for $l = 1,\ 2,\ 3,\ 4$. We note further that the second, $3-2\kappa$-th and fourth row of the matrix $C^{(\kappa)}(\nu-1)A^{*}_{1,0}(1,\nu)$ coincides with respectively the second, $3-2\kappa$-th and fourth row of the matrix $A^*_{1,0}(1,\nu)$. In view of the equalities (148) – (151) in [17], since $\alpha=1$ and $\mu=\mu_1(\nu)=(\nu+1)^2$, $\tau=\tau_1(\nu)=\nu+1$ now, it follows that

$$\nu^5=p_5(\mu)+\tau q_5(\mu), \tag{116}$$

where

$$p_5(\mu)=-5\mu^2-10\mu-1,\ q_5(\mu)=\mu^2+10\mu+5. \tag{117}$$

Let $A^{\vee\kappa\vee}_{1,0}(1;\nu)$ denotes the 4×4-matrix with second, $(3-2\kappa)$-th and fourth row equal respectively to the second, $(3-2\kappa)$-th and fourth row of the matrix $C(\nu-1)A^{\vee}_{1,0}(1,\nu)D(\nu)$ and with $(2\kappa+1)$-th row equal to

$$(-2)^\kappa(\mu_1(\nu)^{2-\kappa})p_5(\mu)\bar{e}_{4,2\kappa+1}.$$

Let further $A^{\wedge\kappa\wedge}_{1,0}(1;\nu)$ denotes the 4×4-matrix with second, $(3-2\kappa)$-th and fourth row equal respectively to the second, $(3-2\kappa)$-th and fourth row of the matrix $C(\nu-1)A^{\wedge}_{1,0}(1,\nu)D(\nu)$ and with $(2\kappa+1)$-th row equal to

$$(-2)^\kappa(\mu_1(\nu)^{2-\kappa})q_5(\mu)\bar{e}_{4,2\kappa+1}.$$

Then, in view of (93), (115) and (116),

$$A^{*\kappa*}_{1,0}(1;\nu)=A^{\vee\kappa\vee}_{1,0}(1;\nu)+\tau A^{\wedge\kappa\wedge}_{1,0}(1;\nu), \tag{118}$$

where $\kappa=0$, 1. We denote by $a^{\vee\kappa\vee}_{1,0,i,j}(1;\nu)$, $a^{\wedge\kappa\wedge}_{1,0,i,j}(1;\nu)$ and $a^{*\kappa*}_{1,0,i,j}(1;\nu)$, where $\kappa=0,1$; $i,j=1,2,3,4$, the expressions, which stand on intersection of i-th row and j-th column in the matrices respectively $A^{*\kappa*}_{1,0}(1;\nu)$, $A^{\vee\kappa\vee}_{1,0}(1;\nu)$ and $A^{\wedge\kappa\wedge}_{1,0}(1;\nu)$. Then, in view of (115),

$$a^{*\kappa*}_{1,0,2\kappa+1,2\kappa+1}(1;\nu)= \tag{119}$$

$$(-2)^\kappa\tau^{4-2\kappa}(\tau-1)^5,$$

$$a^{*\kappa*}_{1,0,2\kappa+1,k}(1;\nu)=0, \tag{120}$$

where $\kappa=0$, 1, $k=3-2\kappa$, 2, 4. In view of (60),

$$a^{\vee}_{1,0,1,1}(1;\nu)=-2\mu-21\mu^2. \tag{121}$$

In view of (64),

$$a^{\wedge}_{1,0,1,1}(1;\nu)=10\mu+17\mu^2. \tag{122}$$

In view of (124), (125), (93) and (118),

$$a^{*}_{1,0,1,1}(1;\nu)= \tag{123}$$

$$17\tau^5-21\tau^4+10\tau^3-2\tau^2)=\tau^2(17\tau^3-21\tau^2+10\tau-2).$$

In view of (68), (103), (104),

$$a^{\vee 0\vee}_{1,0,2,1}(1;\nu)=a^{\vee}_{1,0,2,1}(1;\nu)=-\mu-20\mu^2-12\mu^3. \tag{124}$$

In view of (80), (103), (104),

$$a^{\wedge 0\wedge}_{1,0,2,1}(1;\nu)=a^{\wedge}_{1,0,2,1}(1;\nu)=7\mu+26\mu^2. \tag{125}$$

In view of (124), (125), (93) and (118),

$$\begin{gathered} a^{*0*}_{1,0,2,1}(1;\nu)=a^{*}_{1,0,2,1}(1;\nu)= \\ -(12\tau^6-26\tau^5+20\tau^4-7\tau^3+\tau^2)= \\ -\tau^2(\tau-1)(12\tau^3-14\tau^2+6\tau-1)= \\ -\tau^2(\tau-1)(2\tau-1)(6\tau^2-4\tau+1). \end{gathered} \tag{126}$$

In view of (72), (103), (104),

$$a^{\vee 0\vee}_{1,0,3,1}(1;\nu)=a^{\vee}_{1,0,3,1}(1;\nu)=-\mu-21\mu^2-26\mu^3. \tag{127}$$

In view of (84), (103), (104),

$$\begin{gathered} a^{\wedge 0\wedge}_{1,0,3,1}(1;\nu)=a^{\wedge}_{\alpha,0,3,1}(1;\nu)= \\ 7\mu+33\mu^2+8\mu^3. \end{gathered} \tag{128}$$

In view of (127), (128), (93) and (118),

$$\begin{gathered} a^{*0*}_{1,0,3,1}(1;\nu)=a^{*}_{1,0,3,1}(1;\nu)= \\ 8\tau^7-26\tau^6+33\tau^5-21\tau^4+ \\ \tau^3-\tau^2=\tau^2(\tau-1)(8\tau^4-18\tau^3+15\tau^2-6\tau+1)= \\ \tau^2(\tau-1)^2(8\tau^3-10\tau^2+5\tau-1). \\ \tau^2(\tau-1)^2(2\tau-1)(4\tau^2-3\tau+1). \end{gathered} \tag{129}$$

In view of (76), (103), (104),

$$a^{\vee 0\vee}_{1,0,4,1}(1;\nu)=a^{\vee}_{1,0,4,1}(1;\nu)=-\mu-21\mu^2-34\mu^3-4\mu^4. \tag{130}$$

In view of (88), (103), (104),

$$a^{\wedge 0\wedge}_{\alpha,0,4,1}(1;\nu)=a^{\wedge}_{\alpha,0,4,1}(1;\nu)=7\mu+35\mu^2+18\mu^3. \tag{131}$$

In view of (130), (131), (93) and (118),

$$\begin{gathered} a^{*0*}_{1,0,4,1}(1;\nu)=a^{*}_{1,0,4,1}(1;\nu)= \\ -4\tau^8+18\tau^7-34\tau^6+35\tau^5-21\tau^4+ \\ 7\tau^3-\tau^2=-\tau^2(\tau-1)(4\tau^5-14\tau^4+20\tau^3-15\tau^2+6\tau-1)= \end{gathered} \tag{132}$$

$$-\tau^2(\tau-1)^2(4\tau^4-10\tau^3+10\tau^2-5\tau+1)=$$
$$-\tau^2(\tau-1)^3(4\tau^3-6\tau^2+4\tau-1)=$$
$$-\tau^2(\tau-1)^3(2\tau-1)(2\tau^2-2\tau+1).$$

In view of (62), (103), (104),

$$a^{\vee 1\vee}_{1,0,1,3}(1;\nu)=a^{\vee}_{1,0,1,3}(1;\nu)=-2-12\mu. \tag{133}$$

In view of (66), (103), (104),

$$a^{\wedge 1\wedge}_{1,0,1,3}(1;\nu)=a^{\wedge}_{1,0,1,3}(1;\nu)=10, \tag{134}$$

In view of (133), (134), (93) and (118),

$$a^{*1*}_{1,0,1,3}(1;\nu)=a^{*}_{1,0,1,3}(1;\nu)= \tag{135}$$
$$-12\tau^2+10\tau-2=-2(2\tau-1)(3\tau-1).$$

In view of (70), (103), (104),

$$a^{\vee 1\vee}_{1,0,2,3}(1;\nu)=a^{\vee}_{1,0,2,3}(1;\nu)=-1-14\mu. \tag{136}$$

In view of (82), (103), (104),

$$a^{\wedge 1\wedge}_{1,0,2,3}(1;\nu)=a^{\wedge}_{\alpha,0,2,3}(1;\nu)=7+8\mu, \tag{137}$$

In view of (136), (137), (93) and (118),

$$a^{*1*}_{1,0,2,3}(1;\nu)=a^{*}_{1,0,2,3}(1;\nu)= \tag{138}$$
$$8\tau^3-14\tau^2+7\tau-1=$$
$$(\tau-1)(8\tau^2-6\tau+1)=(\tau-1)(2\tau-1)(4\tau-1).$$

In view of (74),

$$a^{\vee}_{1,0,3,3}(1;\nu)=-1-18\mu-9\mu^2. \tag{139}$$

In view of (86),

$$a^{\wedge}_{1,0,3,3}(1;\nu)=7+20\mu+\mu^2. \tag{140}$$

Hence,

$$a^{*}_{1,0,3,3}(1;\nu)=\tau^5-9\tau^4+20\tau^3-18\tau^2+7\tau-1= \tag{141}$$
$$(\tau-1)(\tau^4-8\tau^3+12\tau^2-6\tau+1)=$$
$$(\tau-1)^2(\tau^3-7\tau^2+5\tau-1).$$

In view of (78), (103), (104),

$$a^{\vee 1\vee}_{1,0,4,3}(1;\nu)=a^{\vee}_{1,0,4,3}(1;\nu)=-1-31\mu-48\mu^2-4\mu^3, \tag{142}$$

In view of (90), (103), (104),

$$a^{\wedge 1\wedge}_{\alpha,0,4,3}(1;\nu)=a^{\wedge}_{\alpha,0,4,3}(1;\nu)=9+53\mu+22\mu^2, \tag{143}$$

In view of (142), (143), (93) and (118),

$$a^{*1*}_{1,0,4,3}(1;\nu)= \tag{144}$$

$$-4\tau^6+22\tau^5-48\tau^4+53\tau^3-31\tau^2+9\tau-1=$$
$$-(\tau-1)(4\tau^5-18\tau^4+30\tau^3-23\tau^2+8\tau-1)=$$
$$-(\tau-1)^2(4\tau^4-14\tau^3+16\tau^2-7\tau+1)=$$
$$-(\tau-1)^3(4\tau^3-10\tau^2+6\tau-1)=-(\tau-1)^3(2\tau-1)(2\tau^2-4\tau+1).$$

In view of (61),

$$a^{\vee}_{1,0,1,2}(1;\nu)=-20\mu-24\mu^2, \tag{145}$$

In view of (65),

$$a^{\wedge}_{1,0,1,2}(1;\nu)=4+36\mu, \tag{146}$$

In view of (69),

$$a^{\vee}_{1,0,2,2}(1;\nu)=-14\mu-41\mu^2. \tag{147}$$

In view of (81),

$$a^{\wedge}_{1,0,2,2}(1;\nu)=2+36\mu+17\mu^2. \tag{148}$$

In view of (73),

$$a^{\vee}_{1,0,3,2}(1;\nu)=-14\mu-58\mu^2-12\mu^3. \tag{149}$$

In view of (85),

$$a^{\wedge}_{1,0,3,2}(1;\nu)=2+40\mu+42\mu^2. \tag{150}$$

In view of (77).

$$a^{\vee}_{1,0,4,2}(1;\nu)=-15\mu-79\mu^2-38\mu^3. \tag{151}$$

In view of (89),

$$a^{\wedge}_{1,0,4,2}(1;\nu)=2+47\mu+75\mu^2+8\mu^3. \tag{152}$$

In view of (63), (67), (71), (83), (75) and (87),

$$a^{\vee}_{1,0,k,4}(1;\nu)=a^{\wedge}_{1,0,k,4}(1;\nu)=0. \tag{153}$$

for $k = 1,\, 2,\, 3$. In view of (79),

$$a^{\vee}_{1,0,4,4}(1;\nu) = -1 - 10\mu - 5\mu^2. \tag{154}$$

In view of (91),

$$a^{\wedge}_{1,0,4,4}(1;\nu) = 5 + 10\mu + \mu^2. \tag{155}$$

In view of (103), (104), (106) and (100) with $\alpha = 1$,

$$\begin{gathered} a^{*\kappa*}_{1,0,k,2+2j}(1;\nu) = -r_{1,0,2+2j}(\nu)a^{*}_{1,0,k,2\kappa+1,}(1;\nu) + \\ r_{1,0,2\kappa+1}(\nu)a^{*}_{1,0,k,2+2j}(1;\nu) = (-2)^{\kappa}\mu^{2-\kappa}a^{*}_{1,0,k,2+2j}(1;\nu), \end{gathered} \tag{156}$$

where $j,\, \kappa = 0,\, 1,\, k = 3 - 2\kappa,\, 2,\, 4$. Therefore, in view of (145) – (155),

$$\begin{gathered} a^{*0*}_{1,0,3,2}(1;\nu) = \\ -\tau^5(12\tau^5 - 42\tau^4 + 58\tau^3 - 40\tau^2 + 14\tau - 2) = \\ -\tau^5(\tau-1)(12\tau^4 - 30\tau^3 + 28\tau^2 - 12\tau + 2) = \\ -\tau^5(\tau-1)^2(12\tau^3 - 18\tau^2 + 10\tau - 2) = \\ -\tau^5(\tau-1)^2(2\tau-1)(6\tau^2 - 6\tau + 2) = \\ -2\tau^5(\tau-1)^2(2\tau-1)(\tau^3 - (\tau-1)^3), \end{gathered} \tag{157}$$

$$\begin{gathered} a^{*1*}_{1,0,1,2}(1;\nu) = -2\tau^2(-24\tau^4 + 36\tau^3 - 20\tau^2 + 4\tau) = \\ 8\tau^3(6\tau^3 - 9\tau^2 + 5\tau - 1) = 8\tau^3(2\tau-1)(3\tau^2 - 3\tau + 1) = \\ 8\tau^3(2\tau-1)(\tau^3 - (\tau-1)^3), \end{gathered} \tag{158}$$

$$\begin{gathered} a^{*\kappa*}_{1,0,2,2}(1;\nu) = (-2)^{\kappa}\tau^{4-2\kappa}(17\tau^5 - 41\tau^4 + 36\tau^3 - 14\tau^2 + 2\tau) = \\ (-2)^{\kappa}\tau^{5-2\kappa}(\tau-1)(17\tau^3 - 24\tau^2 + 12\tau - 2) = \\ (-2)^{\kappa}\tau^{5-2\kappa}(\tau-1)(\tau^3 + 2(2\tau-1)^3), \end{gathered} \tag{159}$$

$$\begin{gathered} a^{*\kappa*}_{1,0,4,2}(1;\nu) = \\ (-2)^{\kappa}\tau^{5-2\kappa}(8\tau^6 - 38\tau^5 + 75\tau^4 - 79\tau^3 + 47\tau^2 - 15\tau + 2) = \\ (-2)^{\kappa}\tau^{5-2\kappa}(\tau-1)(8\tau^5 - 30\tau^4 + 45\tau^3 - 34\tau^2 + 13\tau - 2) = \\ (-2)^{\kappa}\tau^{5-2\kappa}(\tau-1)^2(8\tau^4 - 22\tau^3 + 23\tau^2 - 11\tau + 2) = \\ (-2)^{\kappa}\tau^{5-2\kappa}(\tau-1)^3(8\tau^3 - 14\tau^2 + 9\tau - 2) = \\ (-2)^{\kappa}\tau^{5-2\kappa}(\tau-1)^3(2\tau-1)(4\tau^2 - 5tau + 2). \end{gathered} \tag{160}$$

$$a^{*\kappa*}_{1,0,k,4}(1;\nu) = 0 \text{ for } k = 1,\, 2,\, 3, \tag{161}$$

(162)
$$a^{*\kappa*}_{1,0,4,4}(1;\nu) =$$
$$(-2)^{\kappa}\tau^{4-2\kappa}(\tau^5-5\tau^4+10\tau^3-10\tau^2+5\tau-1) =$$
$$(-2)^{\kappa}\tau^{4-2\kappa}(\tau-1)^5,$$

where $\kappa = 0,\,1$. Let

(163)
$$\theta(k,\kappa) = \det\begin{pmatrix} r_{1,0,1+2\kappa} & r_{1,0,3-2\kappa} \\ a^{*}_{1,0,k,2\kappa+1}(1;\nu) & a^{*}_{1,0,k,3-2\kappa}(1;\nu), \end{pmatrix},$$

where $\kappa = 0,\,1$, $k = 1,\,2,\,3,\,4$. Then

(164)
$$\theta(k,\kappa) = -\theta(k,1-\kappa),\ \text{for } \kappa = 0,\,1,\ k = 1,\,2,\,3,\,4;$$

moreover, in view of (100) with $\alpha = 1$, (103), (104) and (106),

(165)
$$a^{*\kappa*}_{1,0,k,3-2\kappa}(1;\nu) = \theta(k,\kappa) = -r_{1,0,3-2\kappa)}(\nu)a^{*}_{1,0,k,2\kappa+1}(1;\nu) +$$
$$r_{1,0,1+2\kappa}(\nu)a^{*}_{1,0,k,3-2\kappa}(1;\nu) = -(-2)^{1-\kappa}\mu^{1+\kappa}a^{*}_{1,0,k,2\kappa+1}(1;\nu) +$$
$$(-2)^{\kappa}\mu^{2-\kappa}a^{*}_{1,0,k,3-2\kappa}(1;\nu),$$

where $\kappa = 0,\,1$, $k = 3-2\kappa,\,2,\,4$. If we replace κ by $\kappa' = 1-\kappa$ in (165), then we obtain the equality

(166)
$$a^{*\kappa'*}_{1,0,k,3-2\kappa'}(1;\nu) = \theta(k,\kappa')$$

where $\kappa' = 1-\kappa = 0,\,1$, $k = 3-2\kappa' = 2\kappa+1,\,2,\,4$. Hence,

(167)
$$a^{*\kappa*}_{1,0,k,3-2\kappa}(1;\nu) = -a^{*1-\kappa*}_{1,0,k,2\kappa+1}(1;\nu),$$

where $\kappa = 0,\,1$, $k = 2,\,4$. Therefore, in view of (123) – (144),

(168)
$$a^{*1*}_{1,0,1,1}(1;\nu) = -2\tau^4(17\tau^3-21\tau^2+10\tau-2) -$$
$$-2\tau^4(-6\tau^2+5\tau-1) = -2\tau^4(17\tau^3-27\tau^2+15\tau-3) =$$
$$-2\tau^4((\tau-1)^3+2(2\tau-1)^3).$$

(169)
$$a^{*0*}_{1,0,3,3}(1;\nu) =$$
$$\tau^4(\tau-1)^2(\tau^3-7\tau^2+5\tau-1)+$$
$$2\tau^2(\tau^2(\tau-1)^2(8\tau^3-10\tau^2+5\tau-1)) =$$
$$\tau^4(\tau-1)^2(17\tau^3-27\tau^2+15\tau-3) =$$
$$\tau^4(\tau-1)^2((\tau-1)^3+2(2\tau-1)^3) =$$
$$-\frac{1}{2}(\tau-1)^2a^{*1*}_{1,0,1,1}(1;\nu).$$

$$\begin{gathered} a^{*0*}_{1,0,2,3}(1;\nu) = -a^{*1*}_{1,0,2,1}(1;\nu) = \\ \tau^4((\tau-1)(2\tau-1)(4\tau-1))+ \\ 2\tau^2(-\tau^2(\tau-1)(2\tau-1)(6\tau^2-4\tau+1) = \\ -\tau^4(\tau-1)(2\tau-1)(12\tau^2-12\tau+3) = \\ -3\tau^4(\tau-1)(2\tau-1)^3. \end{gathered} \tag{170}$$

$$\begin{gathered} a^{*0*}_{1,0,4,3}(1;\nu) = -a^{*1*}_{1,0,4,1}(1;\nu) = \\ \tau^4(-(\tau-1)^3(2\tau-1)(2\tau^2-4\tau+1))+ \\ 2\tau^2(-\tau^2(\tau-1)^3(2\tau-1)(2\tau^2-2\tau+1)) = \\ -\tau^4(\tau-1)^3(2\tau-1)(6\tau^2-8\tau+3). \end{gathered} \tag{171}$$

§5. Further properties of the matrix $A^{*\kappa*}_{1,0}(z,\nu)$.

In view of (119), (162)

$$\begin{gathered} \lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-9+2\kappa}a^{\vee\kappa\vee}_{1,0,4,4}(1;\nu) = \\ \lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-9+2\kappa}a^{\vee\kappa\vee}_{1,0,2\kappa+1,2\kappa+1}(1;\nu) = 1, \end{gathered} \tag{172}$$

In view of (120)

$$\lim_{\nu\to+\infty}(-2)^{\kappa}\nu^{l}a^{*\kappa*}_{1,0,2\kappa+1,k}(1;\nu) = 0, \tag{173}$$

where $\kappa = 0$, 1, $k = 3-2\kappa$, 2, 4, $l \in \mathbb{Z}$. In view of (126) and (138),

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-6}a^{*\kappa*}_{1,0,2,2\kappa+1}(1;\nu) = -12(1-\kappa, \tag{174}$$

where $\kappa = 0$, 1. In view of (159), (168) (159) and (169),

$$\begin{gathered} \lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-9+2\kappa}a^{\vee\kappa\vee}_{1,0,2,2}(1;\nu) = \\ \lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-9+2\kappa}a^{\vee\kappa\vee}_{1,0,3-2\kappa,3-2\kappa}(1;\nu) = 17, \end{gathered} \tag{175}$$

where $\kappa = 0$, 1. In view of (170),

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-8}a^{*\kappa*}_{1,0,2,3-2\kappa}(1;\nu) = -2^{-\kappa}24, \tag{176}$$

where $\kappa = 0$, 1. In view of (161),

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{l}a^{**}_{1,0,k,4}(1;\nu) = 0, \tag{177}$$

where $\kappa = 0$, 1, $k = 1\,2$, 3, $l \in \mathbb{Z}$. In view of (129), (135),

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-7+4\kappa}a^{*\kappa*}_{1,0,3-2\kappa,1+2\kappa}(1;\nu) = 8(1-\kappa) \tag{178}$$

In view of (157) and (158)

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-10+4\kappa}a^{**}_{1,0,3-2\kappa,2}(1;\nu) = -2^{\kappa}12, \tag{179}$$

where $\kappa = 0$, 1. In view of (132) and (144)

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-8+2\kappa}a^{*\kappa*}_{1,0,4,2\kappa+1}(1;\nu) = -(-2)^{-\kappa}4, \tag{180}$$

where $\kappa = 0$, 1. In view of (160),

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-11+2\kappa}a^{*\kappa*}_{1,0,4,2}(1;\nu) = 8, \tag{181}$$

where $\kappa = 0$, 1. In view of (171),

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-10}a^{*\kappa*}_{1,0,4,3-2\kappa}(1;\nu) = -2^{-\kappa}12, \tag{182}$$

where $\kappa = 0$, 1.

Let λ be a variable. We denote by $T_{n,\lambda}$ the diagonal $n \times n$-matrix, i-th diagonal element of which is equal to λ^{i-1} for $i = 1$, ..., n. We denote by $T^{(i)}_{\nu,\lambda}$ $n \times n$-diagnoal matrix such that its i-th diagonal element is equal to λ and and its other diagonal elements are equal to 1. It follows from (172) – (182) that

(183)

$$\lim_{\nu\to+\infty}(-2)^{-\kappa}\nu^{-9+2\kappa}((T^{(2\kappa+1)}_{4,\nu})^{4-2\kappa}T_{4,\nu})^{-1}A^{*\kappa*}_{1,0}(1;\nu)(T^{(2\kappa+1)}_{4,\nu})^{4-2\kappa}T_{4,\nu}) = A^{(\kappa)},$$

where $\kappa = 0, 1$,

$$A^{(0)} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -12 & 17 & -24 & 0 \\ 8 & -12 & 17 & 0 \\ 4 & 8 & -12 & 1 \end{pmatrix}, A^{(1)} = \begin{pmatrix} 17 & -24 & 0 & 0 \\ -12 & 17 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -6 & 8 & 2 & 1 \end{pmatrix}.$$

Let me to make some comments to the equalities (183. Accordng to (173) and (183

$$\lim_{\nu\to+\infty}\nu^{-9}((T^{(1)}_{4,\nu})^{4-l}T_{4,\nu})^{-1}A^{*\kappa*}_{1,0}(1;\nu)T^{(1)}_{4,\nu})^{4-l}T_{4,\nu}) \tag{184}$$

$$= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 17 & -24 & 0 \\ 0 & -12 & 17 & 0 \\ 04 & 8 & -12 & 1 \end{pmatrix},$$

for any $l \in \mathbb{N}$, and, in particular, for $l = 4$. Let Σ_κ be the matrix, which appears after replacement $2\kappa + 1$-th row in E_4 by the row $\begin{pmatrix} 1 & 0 & -2 & 0 \end{pmatrix}$. Then, in view of (102) with $\alpha = 1$ and (103,

$$T^{-1}_{4,\nu}C^{(\kappa)}(\nu)T_{4,\nu} = (T^{2\kappa+1}_{4,\nu})^{4-2\kappa}\Sigma_\kappa.$$

Further we have

$$(-2)^{-\kappa}\nu^{-9+2}\times$$
$$((T^{(2\kappa+1)}_{4,\nu})^{4-2\kappa}T_{4,\nu})^{-1}A^{*\kappa*}_{1,0}(1,\nu)T_{4,\nu}T^{(2\kappa+1)}_{4,\nu})^{4-2\kappa}=$$
$$(T^{(2\kappa+1)}_{4,\nu})^{2\kappa-4}\times$$
$$(T^{-1}_{4,\nu}C^{(\kappa)}(\nu-1)T_{4,\nu})\times$$
$$T^{-1}_{4,\nu}A^{*}_{1,0}(1,\nu)T_{4,\nu}\times$$
$$(T^{-1}_{4,\nu}(C^{(\kappa)})^{-1}T_{4,\nu})=$$
$$\Sigma_\kappa T^{-1}_{4,\nu}A^{*}_{1,0}(1,\nu)T_{4,\nu}(\Sigma_\kappa)^{-1}.$$

According to the results of section 11.1 in [17],

$$\lim_{\nu\to\infty}T^{-1}_{4,\nu}A^{*}_{1,0}(1,\nu)T_{4,\nu}=A^{\sim}=$$
$$\begin{pmatrix}17&-24&0&0\\-12&17&0&0\\8&-12&1&0\\-4&8&-4&1\end{pmatrix}.$$

Further we have

$$\Sigma_0A^{\sim}(\Sigma_0)^{-1}=\begin{pmatrix}1&0&-2&0\\-12&17&0&0\\8&-12&1&0\\-4&8&-4&1\end{pmatrix}\begin{pmatrix}1&0&2&0\\0&1&0&0\\1&0&1&0\\0&0&0&1\end{pmatrix}=$$
$$\begin{pmatrix}1&0&0&0\\-12&17&-24&0\\8&-12&17&0\\-4&8&-12&1\end{pmatrix}=A^{(0)},$$
$$\Sigma_1A^{\sim}(\Sigma_1)^{-1}=\begin{pmatrix}17&-24&0&0\\-12&17&0&0\\1&0&-2&0\\-4&8&-4&1\end{pmatrix}\begin{pmatrix}1&0&0&0\\0&1&0&0\\\frac{1}{2}&0&-\frac{1}{2}&0\\0&0&0&1\end{pmatrix}=$$
$$\begin{pmatrix}17&-24&0&0\\-12&17&0&0\\0&0&1&0\\-6&8&2&1\end{pmatrix}=A^{(1)}.$$

So, I can consider the equalities (183) as some test of previous calculations.

§6. Further properties of the functions considered in the Introduction.

Let $\alpha \geq 1$. Then the function $t^r(R(\alpha, t, \nu))^2$ (see (54)) is regular at $t = \infty$ for $r = 0$, 1, 2, is regular at $t = \infty$ for $r = 3$, $\alpha \geq 2$ and has a pole of first order at $t = \infty$ for $r = 3$, $\alpha = 1$. So, in the case $r = 0$, 1, 2, $\alpha \geq 1$ we have the equalities

$$Res(t^r(R(\alpha, t, \nu))^2, t = \infty) = 0, \tag{185}$$

$$\lim_{t\to\infty} t^r(R(\alpha, t, \nu))^2 = 0, \tag{186}$$

and in the case $r = 3$, $\alpha = 1$ we have the equalities

$$Res(t^3(R(1, t, \nu))^2, t = \infty) = -1, \tag{187}$$

$$\lim_{t\to\infty} t^3(R(1, t, \nu))^2 = 0. \tag{188}$$

In view of (55),

$$\delta^r f^*_{\alpha,0,2}(z, \nu) = \sum_{t=1}^{+\infty} z^{-t}\frac{(\nu+\alpha)!^2}{\nu!^2}(-t)^r(R(\alpha, t, \nu))^2, \tag{189}$$

where we consider $r = 0$, 1, 2, 3. Expanding $\frac{(\nu+\alpha)!^2}{(\nu!)^2}(-t)^r(R(\alpha, t, \nu))^2$ into partial fractions relatively t, we obtain

$$\frac{(\nu+\alpha)!^2}{(\nu!)^2}(-t)^r(R(\alpha, t; \nu))^2 = \sum_{i=1}^{2}\left(\sum_{k=0}^{\nu+\alpha}\beta^{(r)}_{\alpha,0,i,k,\nu}(t+k)^{-i}\right), \tag{190}$$

where $\nu \in \mathbb{N}_0$, $r = 0$, 1, 2, 3, $\alpha \in \mathbb{N}$,

$$\beta^{(r)}_{\alpha,0,2-j,k,\nu} = \frac{(\nu+\alpha)!^2}{(\nu!)^2} \times \frac{1}{j!}\lim_{t\to -k}\left(\frac{\partial}{\partial t}\right)^j((-t)^r(R(\alpha, t, \nu)(t+k))^2), \tag{191}$$

for $j = 0$, 1. In view of (185), (190) and (187),

$$\sum_{k=0}^{\nu+\alpha}\beta^{(r)}_{\alpha,0,1,k,\nu} = 0, \tag{192}$$

for $r = 0$, 1, 2 and $\alpha \in \mathbb{N}$.

$$\sum_{k=0}^{\nu+1}\beta^{(3)}_{1,0,1,k,\nu} = -(\nu+1)^2. \tag{193}$$

In view of (190),

$$-\frac{(\nu+\alpha)!^2}{(\nu!)^2}\frac{\partial}{\partial t}((-t)^r(R(\alpha,t;\nu))^2)= \tag{194}$$

$$\sum_{i=1}^{2}\left(\sum_{k=0}^{\nu+\alpha}\beta^{(r)}_{\alpha,0,i,k,\nu}i(t+k)^{-i-1}\right),$$

where $\nu\in\mathbb{N}_0$, $r=0,\ 1,\ 2,\ 3$. Let

$$S_{i,k}(\alpha,\nu)=-\left(\sum_{\kappa=k+1}^{\nu+k}1/\kappa^i\right)- \tag{195}$$

$$\left(\sum_{\kappa=1}^{\nu+\alpha-k}1/\kappa^i\right)+\sum_{\kappa=1}^{k}1/\kappa^i,$$

where $\nu\in\mathbb{N}_0$, $i\in\mathbb{N}$, $k\in[0,\ \nu+\alpha\cap\mathbb{Z}$. In particular,

$$S_{1,0}(0,0)=0,\ S_{1,0}(0,1)=-2,\ S_{1,1}(0,1)=\frac{1}{2}, \tag{196}$$

$$S_{1,0}(1,0)=-1,\ S_{1,1}(1,0)=1, \tag{197}$$

$$S_{1,0}(1,1)=-\frac{5}{2},\ S_{1,1}(1,1)=-\frac{1}{2},\ S_{1,2}(1,1)=\frac{7}{6}, \tag{198}$$

$$S_{1,0}(1,2)=-(1+1/2)-(1+1/2+1/3)=-\frac{10}{3} \tag{199}$$

$$S_{1,1}(1,2)=-(1/2+1/3)-(1+1/2)+1=-\frac{4}{3} \tag{200}$$

$$S_{1,2}(1,2)=-(1/3+1/4)-1+(1+1/2)=-\frac{1}{12} \tag{201}$$

$$S_{1,3}(1,2)=-(1/4+1/5)+(1+1/2+1/3)=\frac{83}{60} \tag{202}$$

In view of (191), (54) and (195)

$$\beta^{(0)}_{\alpha,0,2,k,\nu}= \tag{203}$$

$$\left(\frac{(\nu+\alpha)!}{\nu!}\times\frac{(\nu+k)!}{k!}\times\frac{1}{(\nu+\alpha-k)!}\right)^2=\binom{\nu+\alpha}{k}^2\binom{\nu+k}{k}^2,$$

(204) $$\beta^{(0)}_{\alpha,0,1,k,\nu}=2\beta^{(0)}_{\alpha,0,2,k,\nu}S_{1,k}(\alpha,\nu),$$

where $\nu\in\mathbb{N}_0$, $i\in\mathbb{N}$, $k\in[0,\ \nu+\alpha\cap\mathbb{Z}$. In particular,

(205) $$\beta^{(0)}_{0,0,2,0,0}=\beta^{(0)}_{0,0,2,0,1}=1,\ \ \beta^{(0)}_{0,0,2,1,1}=4,$$

(206) $$\beta^{(0)}_{1,0,2,0,0}=\beta^{(0)}_{1,0,2,1,0}=1,$$

(207) $$\beta^{(0)}_{1,0,2,0,1}=1,\ \beta^{(0)}_{1,0,2,1,1}=16,\ \beta^{(0)}_{1,0,2,2,1}=9,$$

(208) $$\beta^{(0)}_{1,0,2,0,2}=1,\ \beta^{(0)}_{1,0,2,1,2}=81,$$

(209) $$\beta^{(0)}_{1,0,2,2,2}=324,\ \beta^{(0)}_{1,0,2,3,2}=100.$$

In view of (204), (205) – (209) and (196) – (202),

(210) $$\beta^{(0)}_{0,0,1,0,0}=2\beta^{(0)}_{0,0,2,0,0}S_{1,0}(0,0)=0,$$

(211) $$\beta^{(0)}_{0,0,1,0,1}=2\beta^{(0)}_{0,0,2,0,1}S_{1,0}(0,1)=2\times1\times(-2)=-4,$$

(212) $$\beta^{(0)}_{0,0,1,1,1}=2\beta^{(0)}_{0,0,2,1,1}S_{1,1}(0,1)=2\times4\times\frac{1}{2}=4,$$

(213) $$\beta^{(0)}_{1,0,1,0,0}=2\beta^{(0)}_{1,0,2,0,0}S_{1,0}(1,0)=2\times1\times(-1)=-2,$$

(214) $$\beta^{(0)}_{1,0,1,1,0}=2\beta^{(0)}_{1,0,2,1,0}S_{1,1}(1,0)=2\times1\times1=2,$$

(215) $$\beta^{(0)}_{1,0,1,0,1}=2\beta^{(0)}_{1,0,2,0,1}S_{1,0}(1,1)=2\times1\times\frac{-5}{2}=-5,$$

(216) $$\beta^{(0)}_{1,0,1,1,1}=2\beta^{(0)}_{1,0,2,1,1}S_{1,1}(1,1)=2\times16\times\frac{-1}{2}=-16,$$

(217) $$\beta^{(0)}_{1,0,1,2,1}=2\beta^{(0)}_{1,0,2,2,1}S_{1,2}(1,1)=2\times9\times\frac{7}{6}=21,$$

(218) $$\beta^{(0)}_{1,0,1,0,2}=2\beta^{(0)}_{1,0,2,0,2}S_{1,0}(1,2)=2\times1\times\frac{-10}{3}=\frac{-20}{3},$$

$$\beta^{(0)}_{1,0,1,1,2} = 2\beta^{(0)}_{1,0,2,1,2}S_{1,1}(1,2) = 2\times 81\times\frac{-4}{3} = -216, \tag{219}$$

$$\beta^{(0)}_{1,0,1,2,2} = 2\beta^{(0)}_{1,0,2,2,2}S_{1,2}(1,2) = 2\times 324\times\frac{-1}{12} = -54, \tag{220}$$

$$\beta^{(0)}_{1,0,1,3,2} = 2\beta^{(0)}_{1,0,2,3,2}S_{1,3}(1,2) = 2\times 100\times\frac{83}{60} = \frac{830}{3}. \tag{221}$$

We put in (190) $r = 0$, and multiply both sides of obtained equality by $(-t)^r$ for $r = 0$, 1, 2, 3. Then we see that

$$-t\frac{(\nu+\alpha)!^2}{(\nu!)^2}(R(\alpha,t;\nu))^2 = \tag{222}$$

$$\sum_{i=1}^{2}\left(\sum_{k=0}^{\nu+\alpha}\frac{\beta^{(0)}_{\alpha,0,i,k,\nu}(-t-k+k)}{(t+k)^i}\right) = \left(\sum_{k=0}^{\nu+\alpha}\frac{k\beta^{(0)}_{\alpha,0,2,k,\nu}}{(t+k)^2}\right) +$$
$$\left(\sum_{k=0}^{\nu+\alpha}\frac{k\beta^{(0)}_{\alpha,0,1,k,\nu}-\beta^{(0)}_{\alpha,0,2,k,\nu}}{t+k}\right) - \sum_{k=0}^{\nu+\alpha}\beta^{(0)}_{\alpha,0,1,k,\nu},$$

$$(-t)^2\frac{(\nu+\alpha)!^2}{(\nu!)^2}(R(\alpha,t;\nu))^2 = \tag{223}$$

$$\sum_{i=1}^{2}\left(\sum_{k=0}^{\nu+\alpha}\frac{\beta^{(0)}_{\alpha,0,i,k,\nu}(t+k-k)^2}{(t+k)^i}\right) = \left(\sum_{k=0}^{\nu+\alpha}\frac{k^2\beta^{(0)}_{\alpha,0,2,k,\nu}}{(t+k)^2}\right) +$$
$$\left(\sum_{k=0}^{\nu+\alpha}\frac{k^2\beta^{(0)}_{\alpha,0,1,k,\nu}-2k\beta^{(0)}_{\alpha,0,2,k,\nu}}{t+k}\right) + \sum_{k=0}^{\nu+\alpha}(\beta^{(0)}_{\alpha,0,2,k,\nu}+(t-k)\beta^{(0)}_{\alpha,0,1,k,\nu}),$$

$$(-t)^3\frac{(\nu+\alpha)!^2}{(\nu!)^2}(R(\alpha,t;\nu))^2 = \tag{224}$$

$$\sum_{i=1}^{2}\left(\sum_{k=0}^{\nu+\alpha}\frac{\beta^{(0)}_{\alpha,0,i,k,\nu}(-t-k+k)^3}{(t+k)^i}\right) = \left(\sum_{k=0}^{\nu+\alpha}\frac{k^3\beta^{(0)}_{\alpha,0,2,k,\nu}}{(t+k)^2}\right) +$$
$$\left(\sum_{k=0}^{\nu+\alpha}\frac{k^3\beta^{(0)}_{\alpha,0,1,k,\nu}-3k^2\beta^{(0)}_{\alpha,0,2,k,\nu}}{t+k}\right) - \left(\sum_{k=0}^{\nu+\alpha}(t-2k)(\beta^{(0)}_{\alpha,0,2,k,\nu}\right) -$$
$$\sum_{k=0}^{\nu+\alpha}(t^2-kt+k^2)\beta^{(0)}_{\alpha,0,1,k,\nu}.$$

The equality (192) with $r = 0$ again follows from (186) with $r = 1$ and (222); moreover, in view of (190) with $r = 1$, and (222),

$$\beta^{(1)}_{\alpha,0,2,k,\nu} = k\beta^{(0)}_{\alpha,0,2,k,\nu}, \tag{225}$$

$$\beta^{(1)}_{\alpha,0,1,k,\nu} = k\beta^{(0)}_{\alpha,0,1,k,\nu} - \beta^{(0)}_{\alpha,0,2,k,\nu} \tag{226}$$

for $k = 0, ..., \nu+\alpha$, $\alpha \in \mathbb{N}$. The equality (192) with $r = 1$ again follows from (186) with $r = 2$, (192) with $r = 0$, (223) and (226); moreover, in view of (190) with $r = 2$, and (223),

$$\beta^{(2)}_{\alpha,0,2,k,\nu} = k^2\beta^{(0)}_{\alpha,0,2,k,\nu}, \tag{227}$$

$$\beta^{(2)}_{\alpha,0,1,k,\nu} = k^2\beta^{(0)}_{\alpha,0,1,k,\nu} - 2k\beta^{(0)}_{\alpha,0,2,k,\nu} \tag{228}$$

for $\alpha \in \mathbb{N}$ and $k = 0, ..., \nu+\alpha$. The equality (192) with $r = 2$ again follows from (188), (192) with both $r \in \{0, 1\}$, (224), (226) and from (228); moreover, in view of (190) with $r = 3$, and (224),

$$\beta^{(3)}_{\alpha,0,2,k,\nu} = k^3\beta^{(0)}_{\alpha,0,2,k,\nu}, \tag{229}$$

$$\beta^{(3)}_{\alpha,0,1,k,\nu} = k^3\beta^{(0)}_{\alpha,0,1,k,\nu} - 3k^2\beta^{(0)}_{\alpha,0,2,k,\nu} \tag{230}$$

for $\alpha \in \mathbb{N}$ and $k = 0, ..., \nu+\alpha$. In view of (55) – (57),

$$\begin{gathered}(\delta^r)f^*_{\alpha,0,3}(z,\nu) = (\log(z))(\delta)^r)f^*_{\alpha,0,2}(z,\nu) + \\ r(\delta)^{r-1}f^*_{\alpha,0,2}(z,\nu) + (\delta)^r f^*_{\alpha,0,4}(z,\nu) = (\log(z))(\delta)^r)f^*_{\alpha,0,2}(z,\nu) + \\ \sum_{t=1}^{+\infty} z^{-t}\frac{(\nu+\alpha)!^2}{\nu!^2}\left(r(-t)^{r-1} - (-t)^r\frac{\partial}{\partial t}\right)R^2(\alpha,t,\nu) = \\ (\log(z))(\delta)^r)f^*_{\alpha,0,2}(z,\nu) - \sum_{t=1}^{+\infty} z^{-t}\frac{(\nu+\alpha)!^2}{\nu!^2}\frac{\partial}{\partial t}((-t)^r R^2)(\alpha,t,\nu).\end{gathered} \tag{231}$$

In view of (190), (194), (189) (231),

$$\begin{gathered}\delta^r f^*_{\alpha,0,2+j}(z,\nu) - j(\log(z))\delta^r f^*_{\alpha,0,2}(z,\nu) = \\ \sum_{i=1}^{2}\left(\sum_{t=1}^{\infty}\left(\sum_{k=0}^{\nu+\alpha}(1-j+ij)\beta^{(r)}_{\alpha,0,i,k,\nu}z^k z^{-t-k}(t+k)^{-i-j}\right)\right) = \\ \sum_{i=1}^{2}\left(\sum_{k=0}^{\nu+\alpha}(1-j+ij)\beta^{(r)}_{\alpha,0,i,k,\nu}z^k\left(\sum_{t=1}^{\infty} z^{-t-k}(t+k)^{-i-j}\right)\right) = \\ \sum_{i=1}^{2}\left(\sum_{k=0}^{\nu}(1-j+ij)\beta^{(r)}_{\alpha,0,i,k,\nu}z^k\left(L_{i+1}(1/z) - \sum_{\tau=1}^{k} z^{-\tau}i(\tau)^{-i-j}\right)\right) = \\ \left(\sum_{i=1}^{2}(1-j+ij)\beta^{*(r)}_{\alpha,0,i}(z;\nu)L_{i+j}(1/z)\right) - \beta^{(r)}_{\alpha,0,3+j}(z;\nu),\end{gathered} \tag{232}$$

where $j = 0, 1, r = 0, 1, 2, 3, |z| > 1, \alpha \in \mathbb{N}$,

$$L_s(1/z) = \sum_{n=1}^{\infty} 1/(z^n n^s),\ \beta^{*(r)}_{\alpha,0,i}(z;\nu) = \sum_{k=0}^{\nu+\alpha} \beta^{(r)}_{\alpha,0,i,k,\nu} z^k, \tag{233}$$

for $s \in \mathbb{Z}$, $i \in \{1, 2\}$, $\nu \in \mathbb{N}_0$,

$$\beta^{*(r)}_{\alpha,0,3+j}(z;\nu) = \tag{234}$$

$$\sum_{i=1}^{2}\left(\sum_{k=0}^{\nu+\alpha}(1-j+ij)\beta^{(r)}_{\alpha,0,i,k,\nu}\left(\sum_{\tau=1}^{k} z^{k-\tau}(\tau)^{-i-j}\right)\right) =$$

$$\sum_{\sigma=0}^{\nu+\alpha-1} z^{\sigma} \sum_{\tau=1}^{\nu+\alpha-\sigma} \sum_{i=1}^{2}(1-j+ij)\beta^{(r)}_{\alpha,0,i,\sigma+\tau,\nu}(\tau)^{-i-j}.$$

In view of (192) and (233), if $r = 0, 1, 2, \alpha \in \mathbb{N}$. then

$$\beta^{*(r)}_{\alpha,0,1}(z;\nu) = (z-1)\beta^{*\vee(r)}_{\alpha,0,1}(z;\nu), \tag{235}$$

where $\beta^{*\vee(r)}_{\alpha,0,1}(z;\nu) \in \mathbb{Q}[z]$, when $\nu \in \mathbb{N}_0$. In view of (193) and (233),

$$\beta^{*(3)}_{1,0,1}(z;\nu) = -(\nu+1)^2 + (z-1)\beta^{*\vee(3)}_{1,0,1}(z;\nu), \tag{236}$$

where $\beta^{*\vee(3)}_{1,0,1}(z;\nu) \in \mathbb{Q}[z]$, when $\nu \in \mathbb{N}_0$. In view of (225) – (230), (233),

$$\beta^{*(1)}_{\alpha,0,2}(z;\nu) = \delta\beta^{*(0)}_{\alpha,0,2}(z;\nu), \tag{237}$$

$$\beta^{*(1)}_{\alpha,0,1}(z;\nu) = \delta\beta^{*(0)}_{\alpha,0,1}(z;\nu) - \beta^{*(0)}_{\alpha,0,2}(z;\nu), \tag{238}$$

$$\beta^{*(2)}_{\alpha,0,2}(z;\nu) = \delta^2\beta^{*(0)}_{\alpha,0,2}(z;\nu), \tag{239}$$

$$\beta^{*(2)}_{\alpha,0,1}(z;\nu) = \delta^2\beta^{*(0)}_{\alpha,0,1}(z;\nu) - 2\delta\beta^{*(0)}_{\alpha,0,2}(z;\nu), \tag{240}$$

$$\beta^{*(3)}_{\alpha,0,2}(z;\nu) = \delta^3\beta^{*(0)}_{\alpha,0,2}(z;\nu), \tag{241}$$

$$\beta^{*(3)}_{\alpha,0,1}(z;\nu) = \delta^3\beta^{*(0)}_{\alpha,0,1}(z;\nu) - 3\delta^2\beta^{*(0)}_{\alpha,0,2}(z;\nu), \tag{242}$$

where $\alpha \in \mathbb{N}$. Clearly,

$$(-\delta)^k L_n(1/z) = L_{n-k}(1/z), \tag{243}$$

where $k \in [0, +\infty) \cap \mathbb{Z}$, $n \in \mathbb{Z}$, $|z| > 1$,

$$L_1(1/z) = -\log(1-1/z),\ -\delta L_1(1/z) = \tag{244}$$

$$L_0(1/z)=\frac{1}{z-1},\ \delta^2L_1(1/z)=$$

$$L_{-1}(1/z)=\frac{1}{z-1}+\frac{1}{(z-1)^2},\ -\delta^3L_1(1/z)=$$

$$L_{-2}(1/z)=\frac{1}{z-1}+\frac{3}{(z-1)^2}+\frac{2}{(z-1)^3}.$$

We apply the operator δ to the equality (232) for $r=0,\,1,\,2,\,\alpha\in\mathbb{N}$. Then, in view of (243), we obtain the eqvality

$$\delta^{r+1}f^*_{\alpha,0,2+j}(z,\nu)-j(\log(z))\delta^{r+1}f^*_{\alpha,0,2}(z,\nu)=$$
$$j\delta^r f^*_{\alpha,0,2}(z,\nu)+$$
$$\left(\sum_{i=1}^{2}((1-j+ij)\delta\beta^{*(r)}_{\alpha,0,i}(z;\nu))L_{i+j}(1/z)\right)-\delta\beta^{(r)}_{\alpha,0,3+j}(z;\nu)-$$
$$\left(\sum_{i=1}^{2}(1-j+ij)\beta^{*(r)}_{\alpha,0,i}(z;\nu)L_{i+j-1}(1/z)\right)=$$
$$j\left(\left(\sum_{i=1}^{2}\beta^{*(r)}_{\alpha,0,i}(z;\nu)L_i(1/z)\right)-\beta^{(r)}_{\alpha,0,3}(z;\nu)\right)+$$
$$\left(\sum_{i=1}^{2}((1-j+ij)\delta\beta^{*(r)}_{\alpha,0,i}(z;\nu))L_{i+j}(1/z)\right)-\delta\beta^{(r)}_{\alpha,0,3+j}(z;\nu)-$$
$$\left(\sum_{i=1}^{2}(1-j+ij)\beta^{*(r)}_{\alpha,0,i}(z;\nu)L_{i+j-1}(1/z)\right).\tag{245}$$

It follws from (245) with $j=0$ that

$$\delta^{r+1}f^*_{\alpha,0,2}(z,\nu)=$$
$$\left(\sum_{i=1}^{2}(\delta\beta^{*(r)}_{\alpha,0,i}(z;\nu))L_i(1/z)-\beta^{*(r)}_{\alpha,0,i}(z;\nu)L_{i-1}(1/z)\right)-\delta\beta^{(r)}_{\alpha,0,3}(z;\nu)=$$
$$(\delta\beta^{*(r)}_{\alpha,0,2}(z;\nu))L_2(1/z)+(\delta\beta^{*(r)}_{\alpha,0,1}(z;\nu)-\beta^{*(r)}_{\alpha,0,2}(z;\nu))L_1(1/z)-$$
$$\delta\beta^{*(r)}_{\alpha,0,3}(z;\nu)-\beta^{*(r)}_{\alpha,0,1}(z;\nu)L_0(1/z).\tag{246}$$

In view of (232) with $j=0$, (246), (235),

$$\beta^{*(r)}_{\alpha,0,2}(z;\nu)=\delta\beta^{*(r-1)}_{\alpha,0,2}(z;\nu))=\delta^r\beta^{*(0)}_{\alpha,0,2}(z;\nu),\tag{247}$$

$$\beta^{*(r)}_{\alpha,0,1}(z;\nu))=\delta\beta^{*(r-1)}_{\alpha,0,1}(z;\nu)-\beta^{*(r-1)}_{\alpha,0,2}(z;\nu)=$$
$$\delta^r\beta^{*(0)}_{\alpha,0,1}(z;\nu)-r\delta^{r-1}\beta^{*(0)}_{\alpha,0,2}(z;\nu),\tag{248}$$

$$\beta^{*(r)}_{\alpha,0,3}(z;\nu) = \delta\beta^{*(r-1)}_{\alpha,0,3}(z;\nu) + \beta^{*(r-1)}_{\alpha,0,1}(z;\nu))L_0(1/z) = \tag{249}$$

$$\delta\beta^{*(r-1)}_{\alpha,0,3}(z;\nu) + \beta^{*\vee(r-1)}_{\alpha,0,1}(z;\nu),$$

where $r = 1, 2, 3$, and $\alpha \in \mathbb{N}$. The equalities (237) – (242) follow from the equalities (247) and (248) again. In view of (203), (233), (53) and (247)

$$\beta^{*(r)}_{\alpha,0,2}(z;\nu) = \delta^r f^*_{\alpha,0,1}(z;\nu) \in \mathbb{N}[z], \tag{250}$$

where $\alpha \in \mathbb{N}$, $\nu \in \mathbb{N}_0$, $r = 0, 1, 2, 3$. It follws from (245) with $j = 1$ that

$$\delta^{r+1} f^*_{\alpha,0,3}(z,\nu) = (\log(z))\delta^{r+1} f^*_{\alpha,0,2}(z,\nu) + \tag{251}$$

$$\left(\left(\sum_{i=1}^{2}\beta^{*(r)}_{\alpha,0,i}(z;\nu)L_i(1/z)\right) - \beta^{(r)}_{\alpha,0,3}(z;\nu)\right) +$$

$$\left(\sum_{i=1}^{2} i(\delta\beta^{*(r)}_{\alpha,0,i}(z;\nu))L_{i+1}(1/z)\right) - \delta\beta^{(r)}_{\alpha,0,4}(z;\nu) -$$

$$\left(\sum_{i=1}^{2} i\beta^{*(r)}_{\alpha,0,i}(z;\nu)L_i(1/z)\right) = (\log(z))\delta^{r+1} f^*_{\alpha,0,2}(z,\nu) +$$

$$\left(\sum_{i=1}^{2} i(\delta\beta^{*(r)}_{\alpha,0,i}(z;\nu))L_{i+1}(1/z)\right) - \delta\beta^{(r)}_{\alpha,0,4}(z;\nu) -$$

$$\beta^{*(r)}_{\alpha,0,2}(z;\nu)L_2(1/z) - (\beta^{(r)}_{\alpha,0,3}(z;\nu) + \delta\beta^{(r)}_{\alpha,0,4}(z;\nu)) =$$

$$(\log(z))\delta^{r+1} f^*_{\alpha,0,2}(z,\nu) + 2(\delta\beta^{*(r)}_{\alpha,0,2}(z;\nu))L_3(1/z) +$$

$$(\delta\beta^{*(r)}_{\alpha,0,1}(z;\nu) - \beta^{*(r)}_{\alpha,0,2}(z;\nu))L_2(1/z) - (\delta\beta^{(r)}_{\alpha,0,4}(z;\nu) + \beta^{*(r)}_{\alpha,0,3}(z;\nu)).$$

In view of (232) with $j = 1$, (251),

$$\beta^{*(r+1)}_{\alpha,0,2}(z;\nu) = \delta\beta^{*(r)}_{\alpha,0,2}(z;\nu) = \delta^{r+1}\beta^{*(0)}_{\alpha,0,2}(z;\nu), \tag{252}$$

$$\beta^{*(r+1)}_{\alpha,0,1}(z;\nu) = \delta\beta^{*(r)}_{\alpha,0,2}(z;\nu) - \beta^{*(r)}_{\alpha,0,2}(z;\nu) = \tag{253}$$

$$\delta^{r+1}\beta^{(0)}_{\alpha,0,1}(z;\nu) - \delta^r\beta^{(0)}_{\alpha,0,2}(z;\nu),$$

where $r = 0, 1, 2$, and we obtain (247) – (247) again. Moreover,

$$\beta^{*(r+1)}_{\alpha,0,4}(z;\nu) = \delta\beta^{*(r)}_{\alpha,0,4}(z;\nu) + \beta^{*(r)}_{\alpha,0,3}(z;\nu), \tag{254}$$

where $r = 0, 1, 2$. If we take now $z \in (1, +\infty)$ and will tend z to 1, then, in view of (232), (235), (236) and (244)

$$\delta^r f^*_{\alpha,0,2+j}(1,\nu) = \lim_{z\to 1+0}\delta^r f^*_{\alpha,0,2}(z,\nu) = \tag{255}$$

$$(1 - i + ij\beta^{*(r)}_{\alpha,0,2}(1;\nu)\zeta(2+j) - \beta^{*(r)}_{\alpha,0,3+j}(1;\nu),$$

where $r = 0, 1, 2$, $j = 0, 1$,

$$\lim_{z\to 1+0}(z-1)\delta^3 f^*_{\alpha,0,2}(z,\nu) = 0, \tag{256}$$

if $\alpha \in \mathbb{N}$. In view of (233), (234), (205) – (217),

$$\beta^{*(0)}_{0,0,1}(z;0) = \beta^{(0)}_{0,0,1,0,0} = 0, \tag{257}$$

$$\beta^{*(0)}_{0,0,2}(z;0) = \beta^{(0)}_{0,0,2,0,0} = 1, \tag{258}$$

$$\beta^{*(0)}_{0,0,3}(z;0) = \beta^{*(0)}_{0,0,4}(z;0) = 0, \tag{259}$$

$$\beta^{*(0)}_{0,0,1}(z;1) = \beta^{(0)}_{0,0,1,0,1} + \beta^{(0)}_{0,0,1,1,1}z = -4 + 4z, \tag{260}$$

$$\beta^{*(0)}_{0,0,2}(z;1) = \beta^{(0)}_{0,0,2,0,1} + \beta^{(0)}_{0,0,2,1,1}z = 1 + 4z, \tag{261}$$

$$\beta^{*(0)}_{0,0,3}(z;1) = \beta^{(0)}_{0,0,1,1,1} + \beta^{(0)}_{0,0,2,1,1} = 8, \tag{262}$$

$$\beta^{*(0)}_{0,0,4}(z;1) = \beta^{(0)}_{0,0,1,1,1} + 2\beta^{(0)}_{0,0,2,1,1} = 12, \tag{263}$$

$$\beta^{*(0)}_{1,0,1}(z;0) = \beta^{(0)}_{1,0,1,0,0} + \beta^{(0)}_{1,0,1,1,0}z = -2 + 2z, \tag{264}$$

$$\beta^{*(0)}_{1,0,2}(z;0) = \beta^{(0)}_{1,0,2,0,0} + \beta^{(0)}_{1,0,2,1,0}z = 1 + z, \tag{265}$$

$$\beta^{*(0)}_{1,0,3}(z;0) = \beta^{(0)}_{1,0,1,1,0} + \beta^{(0)}_{1,0,2,1,0} = 3, \tag{266}$$

$$\beta^{*(0)}_{1,0,4}(z;0) = \beta^{(0)}_{1,0,1,1,0} + 2\beta^{(0)}_{1,0,2,1,0} = 4, \tag{267}$$

$$\beta^{*(0)}_{1,0,1}(z;1) = \beta^{(0)}_{1,0,1,0,1} + \beta^{(0)}_{1,0,1,1,1}z + \\ \beta^{(0)}_{1,0,1,2,1}z^2 = -5 - 16z + 21z^2 = (z-1)(21z+5), \tag{268}$$

$$\beta^{*(0)}_{1,0,2}(z;1) = \beta^{(0)}_{1,0,2,0,1} + \beta^{(0)}_{1,0,2,1,1}z + \\ \beta^{(0)}_{1,0,2,2,1}z^2 = 1 + 16z + 9z^2, \tag{269}$$

$$\beta^{*(0)}_{1,0,3}(z;1) = \beta^{(0)}_{1,0,1,1,1} + \beta^{(0)}_{1,0,2,1,1} + \tag{270}$$

$$\frac{1}{2}\beta^{(0)}_{1,0,1,2,1}+\frac{1}{4}\beta^{(0)}_{1,0,2,2,1}+(\beta^{(0)}_{1,0,1,2,1}+\beta^{(0)}_{1,0,2,2,1})z=$$

$$-16+16+\frac{1}{2}\times 21+\frac{1}{4}\times 9+(21+9)z=\frac{51}{4}+30z,$$

$$\beta^{*(0)}_{1,0,4}(z;1)=\beta^{(0)}_{1,0,1,1,1}+2\beta^{(0)}_{1,0,2,1,1}+ \tag{271}$$

$$\frac{1}{4}\beta^{(0)}_{1,0,1,2,1}+2\times\frac{1}{8}\beta^{(0)}_{1,0,2,2,1}+(\beta^{(0)}_{1,0,1,2,1}+2\beta^{(0)}_{1,0,2,2,1})z=$$

$$-16+2\times 16+\frac{1}{4}\times 21+\frac{1}{4}\times 9+(21+18)z=\frac{47}{2}+39z.$$

$$\beta^{*(0)}_{1,0,1}(z;2)=\beta^{(0)}_{1,0,1,0,2}+\beta^{(0)}_{1,0,1,1,2}z+ \tag{272}$$

$$\beta^{(0)}_{1,0,1,2,2}z^2+\beta^{(0)}_{1,0,1,3,2}z^3=$$

$$-\frac{20}{3}-216z-54z^2+\frac{830}{3}z^3=(z-1)(830z^2+668z+20)/3.$$

$$\beta^{*(0)}_{1,0,2}(z;2)=\beta^{(0)}_{1,0,2,0,2}+\beta^{(0)}_{1,0,2,1,2}z+\beta^{(0)}_{1,0,2,2,2}z^2+ \tag{273}$$

$$\beta^{(0)}_{1,0,2,3,2}z^3=1+81z+324z^2+100z^3.$$

$$\beta^{*(0)}_{1,0,3}(z;2)=\beta^{(0)}_{1,0,1,1,2}+\beta^{(0)}_{1,0,2,1,2}+ \tag{274}$$

$$\frac{1}{2}\beta^{(0)}_{1,0,1,2,2}+\frac{1}{4}\beta^{(0)}_{1,0,2,2,2}+\frac{1}{3}\beta^{(0)}_{1,0,1,3,2}+\frac{1}{9}\beta^{(0)}_{1,0,2,3,2}+$$

$$(\beta^{(0)}_{1,0,1,2,2}+\beta^{(0)}_{1,0,2,2,2})z+\left(\frac{1}{2}\beta^{(0)}_{1,0,1,3,2}+\frac{1}{4}\beta^{(0)}_{1,0,2,3,2}\right)z+$$

$$(\beta^{(0)}_{1,0,1,3,2}+\beta^{(0)}_{1,0,2,3,2})z^2=$$

$$-216+81-54/2+324/4+(830+100)/9+((-54+324-54+415/3+100/4)z+$$

$$(830/3+100)z^2=\frac{67}{3}+\frac{1300}{3}z+\frac{1130}{3}z^2.$$

$$\beta^{*(0)}_{1,0,4}(z;2)=\beta^{(0)}_{1,0,1,1,2}+2\beta^{(0)}_{1,0,2,1,2}+ \tag{275}$$

$$\frac{1}{4}\beta^{(0)}_{1,0,1,2,2}+2\times\frac{1}{8}\beta^{(0)}_{1,0,2,2,2}+$$

$$\frac{1}{9}\beta^{(0)}_{1,0,1,3,2}+2\times\frac{1}{27}\beta^{(0)}_{1,0,2,3,2}+$$

$$(\beta^{(0)}_{1,0,1,2,2}+2\beta^{(0)}_{1,0,2,2,2})z+\left(\frac{1}{4}\beta^{(0)}_{1,0,1,3,2}+2\times\frac{1}{8}\beta^{(0)}_{1,0,2,3,2}\right)z+$$

$$(\beta^{(0)}_{1,0,1,3,2}+2\beta^{(0)}_{1,0,2,3,2})z^2=$$

$$-216+162+(-54+324)/4+(830+200)/27+((-57+648+(830/3+100)/4)z+$$

$$(830/3+200)z^2=\frac{2789}{54}+\frac{4129}{6}z+\frac{1430}{3}z^2.$$

In view of (247) – (249), (254) and (264) – (275),

$$\beta^{*(1)}_{1,0,1}(z;0)=\delta\beta^{*(0)}_{1,0,1}(z;0)-\beta^{*(0)}_{1,0,2}(z;0)=\tag{276}$$

$$2z-1-z=z-1,$$

$$\beta^{*(1)}_{1,0,2}(z;0)=\delta\beta^{*(0)}_{1,0,2}(z;0)=z,\tag{277}$$

$$\beta^{*(1)}_{1,0,3}(z;0)=\delta\beta^{*(0)}_{1,0,3}(z;0)+\beta^{*\vee(0)}_{1,0,1}(z;0)=2\tag{278}$$

$$\beta^{*(1)}_{1,0,4}(z;0)=\delta\beta^{*(0)}_{1,0,4}(z;0)+\beta^{*(0)}_{1,0,3}(z;0)=3,\tag{279}$$

$$\beta^{*(1)}_{1,0,1}(z;1)=\delta\beta^{*(0)}_{1,0,1}(z;1)-\beta^{*(0)}_{1,0,2}(z;1)=-16z+\tag{280}$$

$$42z^2-(1+16z+9z^2)=-1-32z+33z^2=(z-1)(33z+1),$$

$$\beta^{*(1)}_{1,0,2}(z;1)=\delta\beta^{*(0)}_{1,0,2}(z;1)=16z+18z^2,\tag{281}$$

$$\beta^{*(1)}_{1,0,3}(z;1)=\delta\beta^{*(0)}_{1,0,3}(z;1)+\beta^{*\vee(0)}_{1,0,1}(z;1)=\tag{282}$$

$$30z+21z+5=51z+5,$$

$$\beta^{*(1)}_{1,0,4}(z;1)=\delta\beta^{*(0)}_{1,0,4}(z;1)+\beta^{*(0)}_{1,0,3}(z;1)=\tag{283}$$

$$39z+\frac{51}{4}+30z=69z+\frac{51}{4},$$

$$\beta^{*(1)}_{1,0,1}(z;2)=\delta\beta^{*(0)}_{1,0,1}(z;2)-\beta^{*(0)}_{1,0,2}(z;2)=\tag{284}$$

$$-216z-108z^2+830z^3-(1+81z+324z^2+100z^3)=$$
$$-1-297z-432z^2+730z^3=(z-1)(730z^2+298z+1),$$

$$\beta^{*(1)}_{1,0,2}(z;2)=\delta\beta^{*(0)}_{1,0,2}(z;2)=81z+648z^2+300z^3,\tag{285}$$

$$\beta^{*(1)}_{1,0,3}(z;2)=\delta\beta^{*(0)}_{1,0,3}(z;2)+\beta^{*\vee(0)}_{1,0,1}(z;2)=\tag{286}$$

$$\frac{1300}{3}z+\frac{2260}{3}z^2+\frac{830}{3}z^2+\frac{668}{3}z+\frac{20}{3}=$$
$$1030z^2+656z+\frac{20}{3},$$

(287)
$$\beta^{*(1)}_{1,0,4}(z;2)=\delta\beta^{*(0)}_{1,0,4}(z;2)+\beta^{*(0)}_{1,0,3}(z;2)=$$
$$\frac{4129}{6}z+\frac{2860}{3}z^2+\frac{67}{3}+\frac{1300}{3}z+\frac{1130}{3}z^2=$$
$$\frac{67}{3}+\frac{6729}{6}z+1330z^2,$$

In view of (247) – (249), (254) and (276) – (287),

(288)
$$\beta^{*(2)}_{1,0,1}(z;0)=\delta\beta^{*(1)}_{1,0,1}(z;0)-\beta^{*(1)}_{1,0,2}(z;0)=z-z=0,$$

(289)
$$\beta^{*(2)}_{1,0,2}(z;0)=\delta\beta^{*(1)}_{1,0,2}(z;0)=z,$$

(290)
$$\beta^{*(2)}_{1,0,3}(z;0)=\delta\beta^{*(1)}_{1,0,3}(z;0)+\beta^{*\vee(1)}_{1,0,1}(z;0)=1,$$

(291)
$$\beta^{*(2)}_{1,0,4}(z;0)=\delta\beta^{*(1)}_{1,0,4}(z;0)+\beta^{*(1)}_{1,0,3}(z;0)=2$$

(292)
$$\beta^{*(2)}_{1,0,1}(z;1)=\delta\beta^{*(1)}_{1,0,1}(z;1)-\beta^{*(1)}_{1,0,2}(z;1)=$$
$$-32z+66z^2-(16z+18z^2)=-48z+48z^2=48z(z-1).$$

(293)
$$\beta^{*(2)}_{1,0,2}(z;1)=\delta\beta^{*(1)}_{1,0,2}(z;1)=16z+36z^2,$$

(294)
$$\beta^{*(2)}_{1,0,3}(z;1)=\delta\beta^{*(1)}_{1,0,3}(z;1)+\beta^{*\vee(1)}_{1,0,1}(z;1)=$$
$$51z+33z+1=84z+1,$$

(295)
$$\beta^{*(2)}_{1,0,4}(z;1)=\delta\beta^{*(1)}_{1,0,4}(z;1)+\beta^{*(1)}_{1,0,3}(z;1)=$$
$$69z+51z+5=120z+5,$$

(296)
$$\beta^{*(2)}_{1,0,1}(z;2)=\delta\beta^{*(1)}_{1,0,1}(z;2)-\beta^{*(1)}_{1,0,2}(z;2)=$$
$$-297z-864z^2+2190z^3-(81z+648z^2+300z^3)=$$
$$z(-378-1512z+1890z^2)=378(z-1)z(5z+1),$$

(297)
$$\beta^{*(2)}_{1,0,2}(z;2)=\delta\beta^{*(1)}_{1,0,2}(z;2)=81z+1296z^2+900z^3,$$

(298)
$$\beta^{*(0)}_{1,0,3}(z;2)=\delta\beta^{*(1)}_{1,0,3}(z;2)+\beta^{*\vee(1)}_{1,0,1}(z;2)$$
$$2060z^2+656z+730z^2+298z+1=2790z^2+954z+1,$$

(299)
$$\beta^{*(2)}_{1,0,4}(z;2)=\delta\beta^{*(1)}_{1,0,4}(z;2)+\beta^{*(1)}_{1,0,3}(z;2)=$$
$$\frac{6729}{6}z+2660z^2+1030z^2+656z+\frac{20}{3}=3690z^2+\frac{3555}{2}z+\frac{20}{3},$$

§7. End of the proof of theorem A.

Let $y^{(\kappa)}_{1,0,i,k}(z;\nu)$ denotes $i-$th element of the column $Y^{(\kappa)}_{1,0,,k}(z;\nu)$ in (109). Then, in view of (95), (103), (109),

$$y^{(\kappa)}_{1,0,i+1,k}(z,\nu) = \delta^i f^{(\kappa)}_{\alpha,0,k}(z,\nu) \tag{300}$$

for $\kappa = 0, 1, i = 1, 2, 3, k = 1, 2, 3, |z| > 1, \nu \in \mathbb{N}_0$. We denote $v^{*\kappa*}_{1,0,i,j}(\nu)$ the expression, which stands in the matrix $V^{*\kappa*}_{1,0}(\nu)$ in intersection of i-th row and j-th column, where $i = 1, 2, 3, 4, j = 1, 2, 3, 4$. Let

$$D_{\alpha,0}(z,\nu,w) = z(w^2 + w(1-\alpha) - \mu_\alpha)^2 - w^4, \tag{301}$$

In view of (100)

$$\frac{1}{z}D_{\alpha,0}(z,\nu,w) = (1-1/z)w^4 + \sum_{k=0}^{3} r_{\alpha,0,k+1}(\nu)w^k, \tag{302}$$

It follows from general properties of Mejer's functions that

$$D_{\alpha,0}(z,\nu,\delta)f_{\alpha,0,k}(z,\nu) = 0, \tag{303}$$

where $|z| > 1, -3\pi/2 < \arg(z) \le \pi/2, \log(z) = \ln(|z|) + i\arg(z), k = 1, 2, 3$. Therefore, in view of (95), (103), (109),

$$y^{(\kappa)}_{2\kappa+1,0,k}(z,\nu) = -(1-1/z)\delta^4 f_{\alpha,0,k}(z,\nu) \tag{304}$$

where

$$\kappa = 0, 1,, |z| > 1, -3\pi/2 < \arg(z) \le \pi/2,$$
$$\log(z) = \ln(|z|) + i\arg(z), k = 1, 2, 3.$$

In view of (53) – (57), (231),

$$\lim_{z\to1+0}(z-1)\delta^4 f_{1,0,2}(z,\nu) =$$
$$\lim_{z\to1+0}(z-1)\left(O(1)\ln\left(1-\frac{1}{z}\right) + \frac{1}{z-1}\right) = 1, \tag{305}$$

$$\lim_{z\to1+0}(z-1)\delta^4 f_{1,0,k}(z,\nu) = 0, \tag{306}$$

if $k - 2 = \pm1$,

$$\lim_{z\to1+0}(\log(z))\delta^i f_{1,0,k}(z,\nu) = \lim_{z\to1+0}(z-1)\delta^i f_{1,0,k}(z,\nu) = 0, \tag{307}$$

if $i = 0, 1, 2, 3, k = 1, 2, 3$. Hence, if we tend $z \in (1,+\infty)$ to 1, then, in view of (95), (103), we obtain the equalities

$$y^{(\kappa)}_{1,0,2\kappa+1,1}(1,\nu) = y_{1,0,2\kappa+1,3}(1,\nu) = 0, y_{1,0,2\kappa+1,2}(1,\nu) = -1. \tag{308}$$

In view of (111), (119) – (171), (300), (304),

$$
\begin{gathered}
-a^{*\kappa*}_{1,0,i,2\kappa+1}(1;\nu)(1-1/z)\delta^4 f_{1,0,k}(z,\nu)+\\
\left(\sum_{j=1}^{2} a^{*\kappa*}_{1,0,i,j+1-\kappa}(1;\nu)\delta^{j-\kappa} f_{1,0,k}(z,\nu)\right)-\\
(z-1)v^{*\kappa*}_{1,0,i,1}(\nu)(1-1/z)\delta^4 f_{1,0,k}(z,\nu)+\\
(z-1)\sum_{j=1}^{2} v^{**}_{1,0,i+1,j+1-\kappa}(\nu)\delta^{j-\kappa} f_{1,0,k}(z,\nu)=\\
(-2)^\kappa\mu_1(\nu)^{2-\kappa}\nu^5\delta^{i-1} f_{\alpha,0,k}(z,\nu-1),
\end{gathered}
\tag{309}
$$

where $i=3-2\kappa,\ 2,\ k=1,\ 2,\ 3,\ |z|>1,\ -3\pi/2<\arg(z)\le\pi/2$ and ν run over the set $M^*_1=((-\infty,-2]\cup[1,+\infty))\cap\mathbb{Z}$. We tend $z\in(1,+\infty)$ to 1 now and obtain the equalities

$$
\begin{gathered}
a^{*\kappa*}_{1,0,i,1}(1;\nu)(k-1)(k-3)+\\
\left(\sum_{j=1}^{2} a^{*\kappa*}_{1,0,i,j+1-\kappa}(1;\nu)(\delta^{j-\kappa} f_{1,0,k})(1,\nu)\right)=\\
(-2)^\kappa\mu_1(\nu)^{2-\kappa}\nu^5\delta^{i-1} i f_{1,0,k}(1,\nu-1),
\end{gathered}
\tag{310}
$$

where $i=2-\kappa,\ 3-\kappa,\ k=1,\ 2,\ 3$ and $\nu\in M^*_1=((-\infty,-2]\cup[1,+\infty))\cap\mathbb{Z}$. Replacing in (310) $\nu\in M^*_1$ by

$$\nu:=-\nu-2\in M^{**}_1=((-\infty,-3]\cup[0,+\infty))\cap\mathbb{Z},$$

and taking in account the equality (110) we obtain the equalities

$$
\begin{gathered}
a^{*\kappa*}_{1,0,i,1}(1;-\nu-2)(k-1)(k-3)+\\
\left(\sum_{j=1}^{2} a^{*\kappa*}_{1,0,i,j+1-\kappa}(1;-\nu-2)(\delta^{j-\kappa} f_{1,0,k})(1,\nu)\right)=\\
(-2)^\kappa\mu_1(\nu)^{2-\kappa}\nu^5\delta^{i-1} f_{1,0,k}(1,\nu+1),
\end{gathered}
\tag{311}
$$

where $i=2-\kappa,\ 3-\kappa,\ k=1,\ 2,\ 3$ and $\nu\in M^{**}_1=((-\infty,-3]\cup[0,+\infty))\cap\mathbb{Z}$. Let

$$
\vec{w}^{(\kappa)}_{i,j}(\nu)=\begin{pmatrix} a^{**}_{1,0,i+1-\kappa,j+1-\kappa}(1;-\nu-2)\\ (1+(-1)^{i+j})/2\\ a^{**}_{1,0,i+1-\kappa,j+1-\kappa}(1;\nu)\end{pmatrix},
\tag{312}
$$

$$
W^{(\kappa)}_i(\nu)=\begin{pmatrix}\vec{w}_{i,1}(\nu) & \vec{w}_{i,2}(\nu)\end{pmatrix},
\tag{313}
$$

$$
Y^{*\kappa**}_k(\nu)=\begin{pmatrix}(\delta^{1-\kappa} f_{1,0,k})(1,\nu)\\ (\delta^{2-\kappa} f_{1,0,k})(1,\nu)\end{pmatrix},
$$

(314)

$$Y^{*\kappa***}_{i,k}(\nu) = \begin{pmatrix} -(-2)^{\kappa}\mu_1(\nu)^{2-\kappa}(\nu+2)^5\delta^{i-\kappa}f_{1,0,k}(1,\nu+1) \\ (\delta^{i-\kappa}f_{1,0,k})(1,\nu) \\ (-2)^{\kappa}\mu_1(\nu)^{2-\kappa}\nu^5\delta^{i-\kappa}f_{\alpha,0,k}(1,\nu-1) \end{pmatrix},$$

where $\kappa = 0,\ 1,\ k = 1,\ 3,\ i = 1,\ 2,\ \nu \in M_1^{****} = ((-\infty,-3] \cup [1,+\infty)) \cap \mathbb{Z}$. Let further

$$\vec{w}^{(\kappa)}_{i,3}(\nu) = \begin{pmatrix} w^{(\kappa)}_{i,3,1}(\nu) \\ w^{(\kappa)}_{i,3,2}(\nu) \\ w^{(\kappa)}_{i,3,3}(\nu) \end{pmatrix} = [\vec{w}^{(\kappa)}_{i,1}(\nu), \vec{w}^{(\kappa)}_{i,2}(\nu)] \tag{315}$$

is vector product of $\vec{w}^{(\kappa)}_{i,1}(\nu)$ and $\vec{w}^{(\kappa)}_{i,2}(\nu)$, and let $\bar{w}^{(\kappa)}_{3,i}(\nu) = (\vec{w}^{(\kappa)}_{3,i}(\nu))^t$ is row conyugte to the column $\vec{w}(\kappa)_{3,i}(\nu)$. Then for scalar products $(\vec{w}^{(\kappa)}_{i,3}(\nu), \vec{w}^{(\kappa)}_{i,j}(\nu))$ we have the equalities

$$\bar{w}^{(\kappa)}_{i,3}(\nu)\vec{w}^{\kappa}_{i,j}(\nu) = (\vec{w}^{(\kappa)}_{i,3}(\nu), \vec{w}^{(\kappa)}_{i,j}(\nu)) = 0,$$

where $\kappa = 0,\ 1,\ i = 1,\ 2,\ j = 1,\ 2$ and

$$\nu \in M_1^{****} = ((-\infty,-3] \cup [1,+\infty)) \cap \mathbb{Z}.$$

Therefore

$$\bar{w}^{(\kappa)}_{i,3}(\nu)W^{(\kappa)}_{i}(\nu) = \begin{pmatrix} 0 & 0 \end{pmatrix}, \tag{316}$$

where $\kappa = 0,\ 1,\ i = 1,\ 2$ and $\nu \in M_1^{****} = ((-\infty,-3] \cup [1,+\infty)) \cap \mathbb{Z}$.

In view of (310) (311) and (316),

$$\bar{w}(\kappa)_{i,3}(\nu)Y^{*\kappa***}_{i,k}(\nu) = \bar{w}(\kappa)_{i,3}(\nu)W(\kappa)_i(\nu)Y^{*\kappa**}_{k}(\nu) = 0, \tag{317}$$

where $\kappa = 0,\ 1,\ i = 1,\ 2,\ k = 1,\ 3$ and $\nu \in M_1^{****} = ((-\infty,-3]\cup[1,+\infty))\cap\mathbb{Z}$. In view of (315) – (313), (118), and (119) – (171)

$$w^(0)_{1,3,1}(\nu) = a^{*0*}_{1,0,2,3}(1;\nu) = -3\tau^4(\tau-1)(2\tau-1)^3, \tag{318}$$

$$w^(1)_{1,3,1}(\nu) = a^{*1*}_{1,0,1,2}(1;\nu) = \tag{319}$$

$$8\tau^3(2\tau-1)(\tau^3-(\tau-1)^3),$$

$$w^{(0)}_{2,3,1,}(\nu) = -a^{*0*}_{1,0,3,2}(1;\nu) = \tag{320}$$

$$2\tau^5(\tau-1)^2(2\tau-1)(\tau^3-(\tau-1)^3),$$

$$w^(1)_{2,3,1}(\nu) = -a^{*1*}_{1,0,2,1}(1;\nu) = a^{*0*}_{1,0,2,3}(1;\nu) = \tag{321}$$

$$-3\tau^4(\tau-1)(2\tau-1)^3 = w^(0)_{1,3,1}(\nu).$$

Since $\tau_\alpha(-\nu-1-\alpha)=-\tau_\alpha(\nu)$, it follows from (318) – (321) that

(322) $$w^{(0)}_{1,3,3}(\nu)=-a^{*0*}_{1,0,2,3}(1;-\nu-2)=3\tau^4(\tau+1)(2\tau+1)^3,$$

(323) $$w^{(0)}_{1,3,3}(\nu)=-a^{*1*}_{1,0,1,2}(1;-\nu-2)=$$
$$-8\tau^3(2\tau+1)((\tau+1)^3-\tau^3),$$

(324) $$w^{(0)}_{2,3,3}(\nu)=a^{*0*}_{1,0,3,2}(1;-\nu-2)=$$
$$-2\tau^5(\tau+1)^2(2\tau+1)((\tau+1)^3-\tau^3),$$

(325) $$w^{(}1)_{2,3,3}(\nu)=a^{*1*}_{1,0,2,1}(1;-\nu-2)=-a^{*0*}_{1,0,2,3}(1;-\nu-2)=$$
$$3\tau^4(\tau+1)(2\tau+1)^3=w^{(}0)_{1,3,3}(\nu).$$

Further we have

(326) $$w^{(0)}_{1,3,2}(\nu)=$$
$$-\det\begin{pmatrix}a^{*(0)*}_{1,0,2,2}(1;-\nu-2) & a^{*(0)*}_{1,0,2,3}(1;-\nu-2)\\ a^{*0*}_{1,0,2,2}(1;\nu) & a^{*0*}_{1,0,2,3}(1;\nu)\end{pmatrix}=$$
$$a^{**}_{1,0,2,2}(1;\nu)a^{**}_{1,0,2,3}(1;-\nu-2)-a^{**}_{1,0,2,3}(1;\nu)a^{**}_{1,0,2,2}(1;-\nu-2)=$$
$$\tau^5(\tau-1)(\tau^3+2(2\tau-1)^3)(-3\tau^4(\tau+1)(2\tau+1)^3)-$$
$$(-3\tau^4(\tau-1)(2\tau-1)^3)(-\tau^5(\tau+1)(\tau^3+2(2\tau+1)^3))=$$
$$-3\tau^9(\tau^2-1)(\tau^3((2\tau-1)^3+(2\tau+1)^3)+4(4\tau^2-1)^3)=$$
$$-12\tau^9(\tau^2-1)(68\tau^6-45\tau^4+12\tau^2-1),$$

(327) $$w^{(0)}_{3,2,2}(\nu)=$$
$$-\det\begin{pmatrix}a^{**}_{1,0,3,2}(1;-\nu-2) & a^{**}_{1,0,3,3}(1;-\nu-2)\\ a^{**}_{1,0,3,2}(1;\nu) & a^{**}_{1,0,3,3}(1;\nu)\end{pmatrix}=$$
$$a^{**}_{1,0,3,2}(1;\nu)a^{**}_{1,0,3,3}(1;-\nu-2)-a^{**}_{1,0,3,3}(1;\nu)a^{**}_{1,0,3,2}(1;-\nu-2)=$$
$$-2\tau^5(\tau-1)^2(2\tau-1)(\tau^3-(\tau-1)^3)\times$$
$$(-\tau^4(\tau+1)^2((\tau+1)^3+2(2\tau+1)^3)-$$
$$(-2\tau^5(\tau+1)^2(2\tau+1)((\tau+1)^3-\tau^3))\times$$
$$(\tau^4(\tau-1)^2((\tau-1)^3+2(2\tau-1)^3)=$$
$$4\tau^9(\tau^2-1)^2(102\tau^6-68\tau^4+21\tau^2-3).$$

(328) $$w^{(1)}_{1,3,2}(\nu)=$$

$$-\det\begin{pmatrix} a^{*(1)*}_{1,0,1,1}(1;-\nu-2) & a^{*(1)*}_{1,0,1,2}(1;-\nu-2) \\ a^{*1*}_{1,0,1,1}(1;\nu) & a^{*1*}_{1,0,1,2}(1;\nu) \end{pmatrix} =$$

$$a^{**}_{1,0,1,1}(1;\nu)a^{**}_{1,0,1,2}(1;-\nu-2) - a^{**}_{1,0,1,2}(1;\nu)a^{**}_{1,0,1,1}(1;-\nu-2) =$$

$$-2\tau^4(17\tau^3 - 27\tau^2 + 15\tau - 3)8\tau^3(6\tau^3 + 9\tau^2 + 5\tau + 1)-$$

$$2\tau^4(17\tau^3 + 27\tau^2 + 15\tau + 3)8\tau^3(6\tau^3 - 9\tau^3 - 5\tau^2 + 1) =$$

$$-32\tau^7(102\tau^6 - 68\tau^4 + 21\tau^2 - 3),$$

(329) $$w^(1)_{2,3,2}(\nu) =$$

$$-\det\begin{pmatrix} a^{*1*}_{1,0,2,1}(1;-\nu-2) & a^{*1*}_{1,0,2,2}(1;-\nu-2) \\ a^{*1*}_{1,0,2,1}(1;\nu) & a^{*1*}_{1,0,2,2}(1;\nu) \end{pmatrix} =$$

$$\det\begin{pmatrix} a^{*1*}_{1,0,2,2}(1;-\nu-2) & a^{*1*}_{1,0,2,1}(1;-\nu-2) \\ a^{*1*}_{1,0,2,2}(1;\nu) & a^{*1*}_{1,0,2,1}(1;\nu) \end{pmatrix} =$$

$$\det\begin{pmatrix} a^{*1*}_{1,0,2,2}(1;-\nu-2) & -a^{**}_{1,0,2,3}(1;-\nu-2) \\ a^{**}_{1,0,2,2}(1;\nu) & -a^{**}_{1,0,2,3}(1;\nu) \end{pmatrix} = -2\tau^{-2}w^{(0)}_{1,3,2}(\nu).$$

In view of (317), (314), (318), (326), (322),

(330) $$(\tau+1)^4(2\tau-1)^3\delta f_{1,0,k}(1,\nu+1) -$$

$$4\tau(68\tau^6 - 45\tau^4 + 12\tau^2 - 1)\delta f_{1,0,k}(1,\nu)-$$

$$(\tau-1)^4(2\tau+1)^3\delta f_{\alpha,0,k}(1,\nu-1) = 0,$$

where $k = 1$, 3 and $\nu \in M_1^{****} = ((-\infty,-3]\cup[1,+\infty))\cap\mathbb{Z}$. According to the equalities (317), (314), (320), (327), (324),

(331) $$(\tau+1)^3(2\tau-1)(\tau^3 - (\tau-1)^3)\delta^2 f_{1,0,k}(1,\nu+1) -$$

$$2(102\tau^6 - 68\tau^4 + 21\tau^2 - 3)\delta^2 f_{1,0,k}(1,\nu)+$$

$$(\tau-1)^3(2\tau+1)((\tau+1)^3 - \tau^3)\delta^2 f_{\alpha,0,k}(1,\nu-1) = 0.$$

In view of (317), (314), (319), (328), (323),

$$8\tau^3(2\tau-1)(\tau^3 - (\tau-1)^3)(2\tau^2(\tau+1)^5)f_{1,0,k}(1,\nu+1)-$$

$$32\tau^7(102\tau^6 - 68\tau^4 + 21\tau^2 - 3)f_{1,0,k}(1,\nu)+$$

$$(-8\tau^3(2\tau+1)((\tau+1)^3 - \tau^3))(-2\tau^2(\tau-1)^5 f_{1,0,k}(1,\nu-1) = 0,$$

(332) $$(\tau+1)^5(2\tau-1)(\tau^3 - (\tau-1)^3)f_{1,0,k}(1,\nu+1) -$$

$$2\tau^2(102\tau^6 - 68\tau^4 + 21\tau^2 - 3)f_{1,0,k}(1,\nu)+$$

$$(\tau-1)^5(2\tau+1)((\tau+1)^3 - \tau^3))f_{1,0,k}(1,\nu-1) = 0,$$

where $k = 1$, 3 and $\nu \in M_1^{****} = ((-\infty,-3]\cup[1,+\infty))\cap\mathbb{Z}$.

Since $\delta^2 f_{1,0,k}(1,\nu+\varepsilon)=\frac{1}{2}(\tau+\varepsilon)^2 f_{1,0,k}(1,\nu+\varepsilon))$ if $k=1$, 3, $\varepsilon^3=\varepsilon$ we can replace $\delta^2 f_{1,0,k}(1,\nu+\varepsilon)$ by $(1/2)(\tau+\varepsilon)^2 f_{1,0,k}(1,\nu+\varepsilon))$ in (330). Then we come to the equality

$$\frac{1}{2}(\tau+1)^5(2\tau-1)(\tau^3-(\tau-1)^3)f_{1,0,k}(1,\nu+1)-$$

$$\tau^2(102\tau^6-68\tau^4+21\tau^2-3)\delta^2 f_{1,0,k}(1,\nu)+$$

$$\frac{1}{2}(\tau-1)^5(2\tau+1)((\tau+1)^3-\tau^3)\delta^2 f_{\alpha,0,k}(1,\nu-1)=0,$$

this equality is equivalent to the equality (332). The equalities (321), (329) and (325) show that we get in case $\kappa=1$ the same result for the relation between $\delta^2 f^*_{1,0,k}(1,\nu+1)$, $\delta^2 f^*_{1,0,k}(1,\nu)$ and $\delta^2 f_{1,0,k}(1,\nu-1)$ with $k=1$, 3, (relation (331)), as it have been obtained in the case $\kappa=0$.

So, in the case $\kappa=0$ we obtain the same results as in the case $\kappa=1$. Since

$$f_{1,0,k}(1,\nu)=\frac{1}{(\nu+1)^2}f^*_{1,0,k}(1,\nu),$$

it follows from (330) – ((331) that

$$(\tau+1)^2\tau(2\tau-1)^3\delta f^*_{1,0,k}(1,\nu+1)- \tag{333}$$

$$4(68\tau^6-45\tau^4+12\tau^2-1)\delta f^*_{1,0,k}(1,\nu)+$$

$$(\tau-1)^2\tau(2\tau+1)^3\delta f^*_{\alpha,0,k}(1,\nu-1)=0,$$

$$(\tau+1)\tau^2(2\tau-1)(\tau^3-(\tau-1)^3)\delta^2 f^*_{1,0,k}(1,\nu+1)- \tag{334}$$

$$2(102\tau^6-68\tau^4+21\tau^2-3)\delta^2 f^*_{1,0,k}(1,\nu)+$$

$$(\tau-1)\tau^2(2\tau+1)((\tau+1)^3-\tau^3)\delta^2 f^*_{\alpha,0,k}(1,\nu-1)=0,$$

where $k=1,3$ Before to complete the proof of Theorem A, we want to check equalities (333) and (334) for $\nu=1$, $k=3$. If $\nu=1$, then the left part of the equality (333) take the form

$$486\delta f^*_{1,0,k}(1,2)-14716\delta f^*_{1,0,k}(1,1)+250\delta f^*_{1,0,k}(1,0)),$$

where $k=1$, 3; in view of (277) – (299),

$$\beta^{*(1)}_{1,0,2}(1;2)=1029,\ \beta^{*(1)}_{1,0,2}(1;1)=34,\ \beta^{*(1)}_{1,0,2}(1;0)=1,$$

$$\beta^{*(1)}_{1,0,4}(1;2)=\frac{14843}{6},\ \beta^{*(1)}_{1,0,4}(1;1)=\frac{327}{4},\ \beta^{*(1)}_{1,0,4}(1;0)=3,$$

$$486\beta^{*(1)}_{1,0,2}(1;2)-14716\beta^{*(1)}_{1,0,2}(1;1)+250\beta^{*(1)}_{1,0,2}(1;0)=$$

$$2(243\times1029-7358\times34+125)=0,$$

$$486\beta^{*(1)}_{1,0,4}(1;2)-14716\beta^{*(1)}_{1,0,4}(1;1)+250\beta^{*(1)}_{1,0,4}(1;0)=$$

$$81 \times 14843 - 3679 \times 327 + 3 \times 250 = 0;$$

therefore, in view of (232), the equality (333) holds for $\nu = 1$. If $\nu = 1$, then the left part of the equality (334) take the form

$$252\delta^2 f^*_{1,0,k}(1,2) - 11042\delta^2 f^*_{1,0,k}(1,\nu) + 380\delta^2 f^*_{\alpha,0,k}(1,0),$$

in view of (289) – (299),

$$\beta^{*(2)}_{1,0,2}(1;2) = 2277,\ \beta^{*(2)}_{1,0,2}(1;1) = 52,\ \beta^{*(2)}_{1,0,2}(1;0) = 1,$$

$$\beta^{*(2)}_{1,0,4}(1;2) = \frac{32845}{6},\ \beta^{*(2)}_{1,0,4}(1;1) = 125,\ \beta^{*(2)}_{1,0,4}(1;0) = 2,$$

$$252\beta^{*(2)}_{1,0,2}(1;2) - 11042\beta^{*(2)}_{1,0,2}(1;1) + 380\beta^{*(2)}_{1,0,2}(1;0) =$$

$$252 \times 2277 - 11042 \times 52 + 380 \times 1 =$$

$$4(63 \times 2277 - 11042 \times 13 + 95) = 0,$$

$$252\beta^{*(2)}_{1,0,4}(1;2) - 11042\beta^{*(2)}_{1,0,4}(1;1) + 380\beta^{*(2)}_{1,0,4}(1;0) =$$

$$42 \times 32845 - 11042 \times 125 + 380 \times 2 =$$

$$10(21 \times 6569 - 5521 \times 25 + 76) = 0;$$

therefore, in view of (232), the equality (334) holds for $\nu = 1$. Let us consider the equations

$$\begin{gathered}(\tau+1)^2\tau(2\tau-1)^3 x_{nu+1} - \\ 4(68\tau^6 - 45\tau^4 + 12\tau^2 - 1)x_\nu + \\ (\tau-1)^2\tau(2\tau+1)^3 x_{\nu-1} = 0,\end{gathered} \tag{335}$$

$$\begin{gathered}(\tau+1)(2\tau-1)\tau^2(\tau^3 - (\tau-1)^3)x_{nu+1} - \\ 2(102\tau^6 - 68\tau^4 + 21\tau^2 - 3)x_n u + \\ (\tau-1)(2\tau+1)\tau^2((\tau+1)^3 - \tau^3)x_{nu-1} = 0,\end{gathered} \tag{336}$$

where $\nu \in \mathbb{N}_0$, $\tau = \nu + 1$. It follows from (333) that $x_\nu = \delta f^*_{1,0,k}(1,\nu)$ satisfies to the equation (335) for $\nu \in \mathbb{N}$ and fixed $k \in \{1,\ 3\}$. It follows from (334) that $x_\nu = \delta^2 f^*_{1,0,k}(1,\nu)$ satisfies (335) for $\nu \in \mathbb{N}$ and fixed $k \in \{1,\ 3\}$. Both equations (335) and (336) are difference equations of Poincaré type with characteristic polynomial $\lambda^2 - 34\lambda + 1$. Hence, if $\{x_\nu\}_{\nu=1}^{+\infty}$ is a non-zero solution of some of these equations, $\varepsilon \in (0,1)$, then there are $C_1(\varepsilon) > 0$ and $C_2(\varepsilon) > 0$ such that only two possibilities exist:

$$\frac{C_1(\varepsilon)}{\left(1+\sqrt{2}\right)^{4\nu(1+\varepsilon)}} \le |x_\nu| \le \frac{C_2(\varepsilon)}{\left(1+\sqrt{2}\right)^{4\nu(1-\varepsilon)}} \tag{337}$$

for all $\nu \in \mathbb{N}$ or

$$C_1(\varepsilon)\left(1+\sqrt{2}\right)^{4\nu(1-\varepsilon)} \le |x_\nu| \le C_2(\varepsilon)\left(1+\sqrt{2}\right)^{4\nu(1+\varepsilon)}. \tag{338}$$

for all $\nu \in \mathbb{N}$. In view of (250), if $x_\nu = \delta^r f^*_{1,0,1}(1,\nu)$ with $r = 1, 2$, then (337) is impossible. In view of (54) with $\alpha \in \mathbb{N}$, (189) and (232) with $j = 1$,

$$\delta^r f^*_{1,0,3}(1,\nu) = (\nu+1)^2 O(1);$$

hence, if $x_\nu = \delta^r f^*_{1,0,3}(1,\nu)$ with $r = 1, 2$, then (338) is impossible. In view of (255) with $j = 1$ and (250), $x_\nu = \beta^{*(1)}_{1,0,4}(1,\nu)$ and $x_\nu = \beta^{*(2)}_{1,0,4}(1,\nu)$ are solutions for $\nu \in \mathbb{N}$ of equations respectively (335) and (336), moreover (338) take place for these solutions. Hence,

$$\frac{C_1(\varepsilon)/C_2(\varepsilon)}{\left(1+\sqrt{2}\right)^{8\nu(1+\varepsilon)}} \leq \left|2\zeta(3) - \frac{\beta^{*(r)}_{1,0,4}(1,\nu)}{\beta^{*(r)}_{1,0,2}(1,\nu)}\right| \leq \frac{C_2(\varepsilon)/C_1(\varepsilon)}{\left(1+\sqrt{2}\right)^{8\nu(1-\varepsilon)}} \tag{339}$$

for $r = 1$, 2. The equations (335) and (336) are equivalent to the equations respectively

$$x_{\nu+1} = b^{(1)}_{\nu+1} x_\nu + a^{(1)}_{\nu+1} x_{\nu-1}, \tag{340}$$

and

$$x_{\nu+1} = b^{(2)}_{\nu+1} x_\nu + a^{(2)}_{\nu+1} x_{\nu-1}, \tag{341}$$

with

$$b^{(1)}_{\nu+1} = b^{(1)}_\tau = \frac{4(68\tau^6 - 45\tau^4 + 12\tau^2 - 1)}{(\tau+1)^2\tau(2\tau-1)^3}, \tag{342}$$

$$a^{(1)}_{\nu+1} = a^{(1)}_\tau = -\frac{(\tau-1)^2\tau(2\tau+1)^3}{(\tau+1)^2\tau(2\tau-1)^3}, \tag{343}$$

$$b^{(2)}_{\nu+1} = b^{(2)}_\tau = \frac{2(102\tau^6 - 68\tau^4 + 21\tau^2 - 3)}{(\tau+1)(2\tau-1)\tau^2(\tau^3 - (\tau-1)^3)}, \tag{344}$$

$$a^{(2)}_{\nu+1} = a^{(2)}_\tau = -\frac{(\tau-1)(2\tau+1)\tau^2((\tau+1)^3 - \tau^3)}{(\tau+1)(2\tau-1)\tau^2(\tau^3 - (\tau-1)^3)} \tag{345}$$

and $\nu \in \mathbb{N}$. To consider the case $\nu = 0$ also, we let

$$a^{(1)}_1 = \frac{-81}{4},\ b^{(1)}_0 = \beta^{*(1)}_{1,0,4}(1,1) = 3,\ b^{(1)}_1 = 34 = \tag{346}$$

$$\left.\frac{4(68\tau^6-45\tau^4+12\tau^2-1)}{(\tau+1)^2\tau(2\tau-1)^3}\right|_{\tau=1},$$

$$a_1^{(2)}=21,\ b_0^{(2)}=\beta^{*(2)}_{1,0,4}(1,0)=2,\ b_1^{(2)}=52= \tag{347}$$

$$\left.\frac{2(102\tau^6-68\tau^4+21\tau^2-3)}{(\tau+1)(2\tau-1)\tau^2(\tau^3-(\tau-1)^3)}\right|_{\tau=1}.$$

The equation (340) is the eqution (1) with $a_\nu^{(1)}$ and $b_\nu^{(1)}$ in the role of respectively a_ν and b_ν. The equation (341) is the eqution (1) with $a_\nu^{(2)}$ and $b_\nu^{(2)}$ in the role of respectively a_ν and b_ν. Let $x_\nu=Q_{1,\nu}^{(1)}$ and $x_\nu=Q_\nu^{(2)}$, where $\nu\in\mathbb{N}_0$, are solutions of the equations respectively (340) and (341) with initial values respectively $x_0=1$, $x_1=b_1^{(1)}=34$ and $x_0=1$, $x_1=b_1^{(2)}=52$. Then we see that $x_\nu=\beta^{*(r)}_{1,0,2}(1,\nu)$ and $x_\nu=Q_\nu^{(r)}$ with fixed $r\in\{1,\,2\}$ are solutions of the same equation with the same initial values. Therefore

$$Q_\nu^{(r)}=\beta^{*(r)}_{1,0,2}(1,\nu) \tag{348}$$

for $r=1,\,2$ and $\nu\in\mathbb{N}_0$. Let $x_\nu=P_\nu^{(1)}$ and $x_\nu=P_\nu^{(2)}$, where $\nu\in\mathbb{N}_0$, are solutions of the equations respectively (340) and (341) with initial values respectively

$$x_0=b_0^{(1)},\ x_1=b_1^{(1)}b_0^{(1)}+a_1^{(1)}=\frac{327}{4}=\beta^{*(1)}_{1,0,4}(1,1)$$

and

$$x_0=b_0^{(2)},\ x_1=b_1^{(2)}b_0^{(2)}+a_1^{(2)}=125=\beta^{*(2)}_{1,0,4}(1,1).$$

Since $x_\nu=\beta^{*(r)}_{1,0,4}(1,\nu)$ and $x_\nu=P_\nu^{(r)}$ with fixed $r\in\{1,\,2\}$ are solutions of the same equation with the same initial values, it follows that

$$P_\nu^{(r)}=\beta^{*(r)}_{1,0,4}(1,\nu) \tag{349}$$

for $r=1,\,2$ and $\nu\in\mathbb{N}_0$. In view of (339) with $r=1$,

$$\frac{C_1(\varepsilon)/C_2(\varepsilon)}{\left(1+\sqrt{2}\right)^{8\nu(1+\varepsilon)}}\le\left|2\zeta(3)-\frac{P_\nu^{(r)}}{Q_\nu^{(r)}}\right|\le\frac{C_2(\varepsilon)/C_1(\varepsilon)}{\left(1+\sqrt{2}\right)^{8\nu(1-\varepsilon)}}. \tag{350}$$

We put in the equation (1) $\xi_\nu=x_\nu/d_\nu$ with some $d_\nu\ne0$ for $\nu\in\mathbb{N}_0$, and we put $d_0=d_{-1}=1$. Then we obtain equation

$$\xi_{\nu+1}=\beta^{\vee}_{\nu+1}\xi_\nu+\alpha_{\nu+1}\xi_{\nu-1} \tag{351}$$

with $\xi_0=x_0=b_0$,

$$\alpha_{\nu+1}=a_{\nu+1}\frac{d_{\nu+1}}{d_{\nu-1}},\beta_{\nu+1}=b_{\nu+1}\frac{d_{\nu+1}}{d_\nu} \tag{352}$$

where $\nu \in \mathbb{N}_0$. Clearly, continuous fraction connected with the equation (351) has the same convergents as continuous fraction corresponding to the equation (1). We apply this transformatin for the equations (340) and (341) instead of equation (1) with respectively

$$d_\nu = d_\nu^{(1)} = \prod_{k=1}^{\nu} k(k+1)^2(2k-1)^3,$$

$$d_\nu = d_\nu^{(2)} = \prod_{k=1}^{\nu} (k+1)(2k-1)k^2(k^3-(k-1)^3),$$

Then we have

$$\frac{d_{\nu+1}^{(1)}}{d_\nu^{(1)}} = (\nu+1)(\nu+2)^2(2\nu+1)^3 = \tau(\tau+1)^2(2\tau-1)^3,$$

$$\frac{d_{\nu+1}^{(2)}}{d_\nu^{(2)}} = (\nu+2)(2\nu+1)(\nu+1)^2((\nu+1)^3-\nu^3) =$$

$$(\tau+1)(2\tau-1)\tau^2(\tau^3-(\tau-1)^3),$$

if $\nu \in \mathbb{N}_0$,

$$\frac{d_{\nu+1}^{(1)}}{d_{\nu-1}^{(1)}} = \tau(\tau+1)^2(2\tau-1)^3 \times (\tau-1)\tau^2(2\tau-3)^3,$$

$$\frac{d_{\nu+1}^{(2)}}{d_{\nu-1}^{(2)}} = (\tau+1)(2\tau-1)\tau^2(\tau^3-(\tau-1)^3)\times$$

$$\tau(2\tau-3)(\tau-1)^2((\tau-1)^3-(\tau-2)^3),$$

if $\nu \in \mathbb{N}$,

$$\frac{d_1^{(1)}}{d_{-1}^{(1)}} = 4, \frac{d_1^{(2)}}{d_{-1}^{(2)}} = 2,$$

and we obtain two expansions of the number $2\zeta(3)$ in continuous fraction: one expansion with $a_1 = -81$, $b_0 = 3$,

$$a_{\nu+1} = a_\tau = -\frac{(\tau-1)^2\tau(2\tau+1)^3}{(\tau+1)^2\tau(2\tau-1)^3}\times$$

$$\tau(\tau+1)^2(2\tau-1)^3 \times (\tau-1)\tau^2(2\tau-3)^3 =$$

$$-(\tau(\tau+1)(4\tau^2-4\tau-3))^3$$

for $\nu \in \mathbb{N}$,

$$b_{\nu+1} = b_\tau = 4(68\tau^6 - 45\tau^4 + 12\tau^2 - 1)$$

for $\nu \in \mathbb{N}_0$, and another expansion with

$$a_1 = 42, \; b_0 = 2, \; a_{\nu+1} = a_\tau =$$

$$-\frac{(\tau-1)(2\tau+1)\tau^2((\tau+1)^3-\tau^3)}{(\tau+1)(2\tau-1)\tau^2(\tau^3-(\tau-1)^3)}\times$$
$$(\tau+1)(2\tau-1)\tau^2(\tau^3-(\tau-1)^3)\times$$
$$\tau(2\tau-3)(\tau-1)^2((\tau-1)^3-(\tau-2)^3)=$$
$$-(\tau-1)^3\tau^3(4\tau^2-4\tau-3)\times$$
$$((\tau+1)^3-\tau^3)((\tau-1)^3-(\tau-2)^3)$$

for $\nu\in\mathbb{N}$,

$$b_{\nu+1}=b_\tau=2(102\tau^6-68\tau^4+21\tau^2-3)$$

for $\nu\in\mathbb{N}_0$, ■

References.

[1] R.Apéry, Interpolation des fractions continues
et irrationalite de certaines constantes,
Bulletin de la section des sciences du C.T.H., 1981, No 3, 37 – 53;
Oskar Perron. Die Lehre von den Kettenbrüche. Dritte, verbesserte und erweiterte Auflage. 1954 B.G.Teubner Verlaggesellshaft. Stuttgart.
Yu.V. Nesterenko. A Few Remarks on $\zeta(3)$,
Mathematical Notes, Vol 59, No 6, 1996,
Matematicheskie Zametki, Vol 59, No 6, pp. 865 – 880 1996
(in Russian).
[4] L.A.Gutnik, On linear forms with coefficients in $\mathbb{N}\zeta(1+\mathbb{N})$
(the detailed version,part 3), Max-Plank-Institut für Mathematik,
Bonn, Preprint Series, 2002, 57, 1 – 33;
[5] ———————, On the measure of nondiscreteness of some modules,
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2005, 32, 1 – 51. *
[6] ———————, On the Diophantine approximations
of logarithms in cylotomic fields.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 147, 1 – 36.
[7] ———————, On some systems of difference equations. Part 1.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 23, 1 – 37.
[8] ———————, On some systems of difference equations. Part 2.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 49, 1 – 31.
[9] ———————, On some systems of difference equations. Part 3.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 91, 1 – 52.
[10] ———————, On some systems of difference equations. Part 4.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 101, 1 – 49.
[11] ———————, On some systems of difference equations. Part 5.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2006, 115, 1 – 9.
[12] ———————, On some systems of difference equations. Part 6.
Max-Plank-Institut für Mathematik, Bonn,
Preprint Series, 2007, 16, 1 – 30.
[13] ———————, On some systems of difference equations. Part 7.
Max-Plank-Institut für Mathematik, Bonn,

Preprint Series, 2007, 53, 1 − 40.
[14] ____________, On some systems of difference equations. Part 8. Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2007, 64, 1 − 44.
[15] ____________, On some systems of difference equations. Part 9. Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2007, 129, 1 − 36.
[16] ____________, On some systems of difference equations. Part 10. Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2007, 131 , 1 − 33.
[17] ____________, On some systems of difference equations. Part 11. Max-Plank-Institut für Mathematik, Bonn, Preprint Series, 2008, 38 , 1 − 45.

E-mail: gutnik@gutnik.mccme.ru